

Connecting via Winsock to STN

18/620,520

Welcome to STN International! Enter x:x

LOGINID:sssptal600txm

PASSWORD:

***** RECONNECTED TO STN INTERNATIONAL *****
SESSION RESUMED IN FILE 'REGISTRY' AT 09:08:22 ON 13 MAY 2006
FILE 'REGISTRY' ENTERED AT 09:08:22 ON 13 MAY 2006
COPYRIGHT (C) 2006 American Chemical Society (ACS)

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	3.52	3.73

=> file reg

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	3.96	4.17

FILE 'REGISTRY' ENTERED AT 09:08:57 ON 13 MAY 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 11 MAY 2006 HIGHEST RN 883943-03-1
DICTIONARY FILE UPDATES: 11 MAY 2006 HIGHEST RN 883943-03-1

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 6, 2006

Please note that search-term pricing does apply when conducting SmartSELECT searches.

*
* The CA roles and document type information have been removed from *
* the IDE default display format and the ED field has been added, *
* effective March 20, 2005. A new display format, IDERL, is now *
* available and contains the CA role and document type information. *
*

Structure search iteration limits have been increased. See HELP SLIMITS for details.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

=>

Uploading C:\Program Files\Stnexp\Queries\09591177b.str

L5 STRUCTURE UPLOADED

=> s l5 sss sam

SAMPLE SEARCH INITIATED 09:09:23 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 21 TO ITERATE

100.0% PROCESSED 21 ITERATIONS 2 ANSWERS
SEARCH TIME: 00.00.01


FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 146 TO 694
PROJECTED ANSWERS: 2 TO 124

L6 2 SEA SSS SAM L5

=> d scan l6

L6 2 ANSWERS REGISTRY COPYRIGHT 2006 ACS on STN
IN Adenosine 5'-(tetrahydrogen triphosphate), 8-(ethylthio)- (9CI)
MF C12 H20 N5 O13 P3 S
CI COM

Absolute stereochemistry.







PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L6 2 ANSWERS REGISTRY COPYRIGHT 2006 ACS on STN
IN Thymidine, 2'-deoxyadenylyl-(3'→5')-2'-deoxyguanylyl-(3'→5')-
2'-deoxycytidylyl-(3'→5')-2'-deoxy-8-(propylthio)adenylyl-
(3'→5')-2'-deoxyguanylyl-(3'→5')-2'-deoxyadenylyl-
(3'→5')- (9CI)
MF C72 H92 N30 O38 P6 S

Absolute stereochemistry.





PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

ALL ANSWERS HAVE BEEN SCANNED

=> s l5 sss full

FULL SEARCH INITIATED 09:10:13 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 351 TO ITERATE

100.0% PROCESSED 351 ITERATIONS

32 ANSWERS

SEARCH TIME: 00.00.01

L7 32 SEA SSS FUL L5

=> file caplus

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

167.38

171.55

FILE 'CAPLUS' ENTERED AT 09:10:18 ON 13 MAY 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 13 May 2006 VOL 144 ISS 21

FILE LAST UPDATED: 11 May 2006 (20060511/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply.
They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> s 17

L8 17 L7

=> d bib abs hitstr 1-17 18

L8 ANSWER 1 OF 17 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2004:911398 CAPLUS

DN 142:214281

TI C8-substituted purine nucleotide analogs and their use as inhibitors of nucleoside triphosphate diphosphohydrolases

IN Halbfinger, Efrat; Fischer, Bilha; Beaudoin, Adrien R.; Gendron, Fernand Pierre

PA Universite de Sherbrooke, Can.; Bar-Ilan University

SO Can. Pat. Appl., 54 pp.

CODEN: CPXXEB

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	CA 2311084	AA	20011209	CA 2000-2311084	20000609

PRAI CA 2000-2311084

OS CASREACT 142:214281

AB Ectonucleoside triphosphate diphosphohydrolases [NTPDases; EC 3.6.1.5] constitute a family of enzymes which are involved in the metabolism of extracellular nucleotides, catalyzing the hydrolysis of the gamma and beta phosphate bonds of triphospho- and diphosphonucleosides (whereas 5'-nucleotidases [EC 3.1.3.5] catalyze the hydrolysis of alpha phosphate bond of monophosphonucleosides). These extracellular nucleotides interact with endothelial, epithelial and smooth muscle cells, as well as blood cells and lymphoid cells, to influence the different physiol. systems of vertebrates. Since these ecto-nucleotidases alter the extracellular concns. of nucleotides these enzymes modulate their physiol. effects, including, for example, platelet aggregation, heart function, control of vascular tone and inflammation reactions, electrolyte secretion and gastrointestinal motility, neurotransmission both in central and peripheral nervous systems, as well as other effects in other physiol. systems. This invention provides C8 substituted purine nucleotide analogs, such as ATP analogs, and further provides their use as inhibitors of NTPDases and thereby as tools to modulate the conversion of nucleotides into nucleoside derivs., and thus modulate the levels of these compds. Such modulation further provides for the modulation of the activity and function of many processes which are affected by these compds.

IT 81609-35-0P 284040-51-3P 284040-52-4P
284040-53-5P


RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(C8-substituted purine nucleotide analogs and their use as inhibitors of nucleoside triphosphate diphosphohydrolases to modulate purine nucleotide levels and biol. processes)

RN 81609-35-0 CAPLUS


CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(ethylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 284040-51-3 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(cycloheptylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 284040-52-4 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-[(2,2-dimethylpropyl)thio]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 284040-53-5 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(hexylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 284040-54-6

RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(C8-substituted purine nucleotide analogs and their use as inhibitors of nucleoside triphosphate diphosphohydrolases to modulate purine nucleotide levels and biol. processes)

RN 284040-54-6 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L8 ANSWER 2 OF 17 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2004:617808 CAPLUS

DN 141:270997

TI Molecular Recognition in Purinergic Receptors. 1. A Comprehensive Computational Study of the h-P2Y1-Receptor

AU Major, Dan T.; Fischer, Bilha

CS Gonda-Goldschmied Medical Research Center, Department of Chemistry, Bar-Ilan University, Ramat-Gan, 52900, Israel

SO Journal of Medicinal Chemistry (2004), 47(18), 4391-4404

CODEN: JMCMAR; ISSN: 0022-2623

PB American Chemical Society

DT Journal

LA English

AB P2Y receptors (P2Y-Rs) are attractive pharmaceutical targets due to their involvement in the modulation of many tissues and organs. The lack of exptl. structural data on P2Y-Rs impedes structure-based drug design. The need to elucidate the receptor's mol. recognition, together with the limitations of previous receptor models, triggered the construction of a new mol. model for the h-P2Y1-R. Therefore, a h-P2Y1-R model was constructed by homol. modeling using the 2.6 Å crystal structure of bovine rhodopsin as a template and subsequently refined by constrained mol. dynamics (MD) simulations in a fully hydrated lipid bilayer environment. ATP was docked into the receptor binding site, followed by binding site refinement using Monte Carlo and MD simulations. Anal. of the h-P2Y1-R-ATP complex suggests that the triphosphate moiety is tightly bound by a multitude of interactions possibly including a Mg²⁺ ion, the ribose ring is not involved in specific interactions, and the adenine ring is bound via N1, N7, and N6. The mol. recognition of the h-P2Y1-R was further probed by ATP derivs. modified on the adenine ring, and correlated with EC50 values for these derivs. Anal. of receptor:ligand complexes and

quantum mech. studies on model compds. support the role of both steric and electronic effects in improving H-bonding (via N1 and N6) and π -stacking interactions. The computed h-P2Y1-R model was validated with respect to our previous biochem. results. The authors believe that this new model of the h-P2Y1-R provides the means for understanding phenomena such as the ligand's potency and receptor subtype selectivity.


IT 284040-54-6

RL: PAC (Pharmacological activity); BIOL (Biological study)
(mol. recognition in human purinergic P2Y1 receptor)

RN 284040-54-6 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 71 THERE ARE 71 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 3 OF 17 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2003:856092 CAPLUS

DN 139:333119

TI Ecto-nucleoside triphosphate diphosphohydrolase inhibition-based methods for screening for a compound useful in the treatment or prevention of lymphocytic disorders, for inhibiting lymphocyte activity and preventing or treating lymphocytic disorders

IN Beaudoin, Adrien; Benrezzak, Ouhida

PA Bioflash Inc., Can.

SO PCT Int. Appl., 63 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003089664	A1	20031030	WO 2003-CA583	20030422
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CA 2382768	AA	20031019	CA 2002-2382768	20020419
	CA 2479501	AA	20031030	CA 2003-2479501	20030422
	AU 2003226989	A1	20031103	AU 2003-226989	20030422
	US 2005164306	A1	20050728	US 2003-511133	20030422
PRAI	CA 2002-2382768	A	20020419		
	WO 2003-CA583	W	20030422		

AB The invention discloses a method of screening for a compound useful in the treatment of a disease or condition characterized by an immune cell disorder, wherein the cell expresses ecto-nucleoside triphosphate diphosphohydrolases (NTPDases), the method comprising contacting a candidate compound with NTPDase, wherein the candidate compound is selected if

the activity of the NTPDase is reduced in the presence of the candidate compound as compared to that in the absence thereof. The invention also discloses a method for inhibiting an immune cell activity in a mammal, comprising targeting immune cells with an effective amount of a NTPDase inhibitor. The invention further discloses a method to prevent or reduce the risk of rejection of transplanted tissue or organ, comprising administering to the animal an effective amount of NTPDase inhibitor.

IT 284040-54-6 344402-39-7


RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(ecto-nucleoside triphosphate diphosphohydrolase inhibition-based methods for screening for agents for treatment of immune cell disorder-associated conditions)

RN 284040-54-6 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 344402-39-7 CAPLUS

CN 5'-Adenylic acid, 8-(butylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 4 OF 17 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2003:753685 CAPLUS

DN 140:205747

TI Characterization and Elucidation of Coordination Requirements of Adenine Nucleotides Complexes with Fe(II) Ions

AU Richter, Yael; Fischer, Bilha

CS Gonda-Goldschmied Medical Research Center, Department of Chemistry, Bar-Ilan University, Ramat-Gan, Israel

SO Nucleosides, Nucleotides & Nucleic Acids (2003), 22(9), 1757-1780
CODEN: NNNAFY; ISSN: 1525-7770

PB Marcel Dekker, Inc.

DT Journal

LA English

AB In spite of the significant role of iron ions-nucleotide complexes in living cells, these complexes have been studied only to a limited extent. Therefore, we fully characterized the ATP:Fe(II) complex including

stoichiometry, geometry, stability consts., and dependence of Fe(II)-coordination on pH. A 1:1 stoichiometry was established for the ATP:Fe(II) complex based on volumetric titrns., UV and SEM/EDX measurements. The coordination sites of ferrous ions in the complex with ATP, established by ¹H-, ³¹P-, and ¹⁵N-NMR, involve the adenine N7 as well as P_α, P_β, and P_γ. Coordination sites remain the same within the pH range of 3.1-8.3. By applying fluorescence monitored Fe(II)-titration, we established a log K value of 5.13 for the Fe(ATP)₂-complex, and 2.31 for the Fe(HATP)- complex. Ferrous complexes of ADP₃- and AMP₂- were less stable (log K 4.43 and 1.68, resp.). The proposed major structure for the Fe(ATP)₂- complex is the open' structure. In the minor closed' structure N7 nitrogen is probably coordinated with Fe(II) through a bridging water mol. The electronic and stereochem. requirements for Fe(II)-coordination with ATP₄- were probed using a series of modified-phosphate or modified-adenine ATP analogs. Fe(II) coordinates solely with the phosphate-oxygen atom, and not with sulfur, amine, or borane in the cases of phosphate-modified analogs of ATP. A high electron d. on N7 and an anti conformation of the adenine-nucleotide are required for enhanced stability of ATP analogs:Fe(II) complexes as compared to ATP complexes (up to more than 100-fold). There are no stereochem. preferences for Fe(II)-coordination with either Rp or Sp isomers of ATP-α-S or ATP-α-BH₃ analogs.

IT 344402-39-7D, iron aquo complexes


RL: FMU (Formation, unclassified); PRP (Properties); FORM (Formation, nonpreparative)

(iron(2+) coordination with adenine nucleotides)

RN 344402-39-7 CAPLUS

CN 5'-Adenylic acid, 8-(butylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 284040-53-5 344402-39-7


RL: RCT (Reactant); RACT (Reactant or reagent)

(iron(2+) coordination with adenine nucleotides)

RN 284040-53-5 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(hexylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 344402-39-7 CAPLUS

CN 5'-Adenylic acid, 8-(butylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RE.CNT 57 THERE ARE 57 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 5 OF 17 CAPLUS COPYRIGHT 2006 ACS on STN
AN 2003:711173 CAPLUS
DN 139:230955
TI Preparation of C8-substituted purine nucleotide analogs as NTPDase
inhibitors
IN Beaudoin, Adrien R.; Gendron, Fernand-Pierre; Halbfinger, Efrat; Fischer,
Bilha
PA Bar-Ilan University, Israel; Universite De Sherbrooke
SO U.S., 21 pp.
CODEN: USXXAM
DT Patent
LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 6617439	B1	20030909	US 2000-591177	20000609
	US 2004043955	A1	20040304	US 2003-620520	20030716
PRAI	US 2000-591177	A3	20000609		
OS	MARPAT 139:230955				
GI					



AB C8-substituted purine nucleotide analogs, I (R is alkyl, cycloalkyl) such as ATP analogs, and their use is described, including their use as inhibitors of NTPDases and thereby as tools to modulate the conversion of nucleotides into nucleoside derivs., and thus modulate the levels of these compds. Such modulation further provides for the modulation of the activity and function of many processes which are affected by these compds. Thus, I [R = (CH2)3Me] was prepared and tested in vivo as NTPDase inhibitor. I [R = (CH2)3Me] interacts specifically with the binding site of the enzyme potentially reduces the risk of interference with other ATP-binding enzymes or receptors, and thus possesses a high degree of specificity. The compds. of the invention were analyzed with resp. to any effects on the activity of purinoceptors.

IT 81609-35-0P 284040-51-3P 284040-52-4P

284040-53-5P 284040-54-6P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of C8-substituted purine nucleotide analogs as NTPDase inhibitors)

RN 81609-35-0 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(ethylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-51-3 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(cycloheptylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-52-4 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-[(2,2-dimethylpropyl)thio]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 284040-53-5 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(hexylthio)- (9CI) (CA INDEX NAME)

NAME)


Absolute stereochemistry.



RN 284040-54-6 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 71 THERE ARE 71 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 6 OF 17 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2001:327844 CAPLUS

DN 135:149038

TI Inhibitors of NTPDase: key players in the metabolism of extracellular purines

AU Gendron, F. P.; Halbfinger, E.; Fischer, B.; Beaudoin, A. R.

CS Department of Biology, University of Sherbrooke, Sherbrooke, Can.

SO Advances in Experimental Medicine and Biology (2000), 486(Purine and Pyrimidine Metabolism in Man X), 119-123

CODEN: AEMBAP; ISSN: 0065-2598

PB Kluwer Academic/Plenum Publishers

DT Journal

LA English

AB This study described the potential of a new class of ATP analogs as nucleoside triphosphate diphosphohydrolase (NTPDase) inhibitors. From previous studies, 8-thiobutyladenosine 5'-triphosphate (8-BuS-ATP) appears to be a specific and efficient NTPDase inhibitor. This novel inhibitor is a new tool to regulate NTPDase activity and thereby influencing purine signaling in mammalian.

IT 81609-35-0 284040-51-3 284040-53-5
284040-54-6

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)


(inhibitors of nucleoside triphosphate diphosphohydrolase - key players in metabolism of extracellular purines)

RN 81609-35-0 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(ethylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

Amkd




RN 284040-51-3 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(cycloheptylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 284040-53-5 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(hexylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 284040-54-6 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 7 OF 17 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2001:293376 CAPLUS

DN 135:41096

TI Novel modified adenosine 5'-triphosphate analogues pharmacologically characterized in human embryonic kidney 293 cells highly expressing rat brain P2Y1 receptor: biotinylated analogue potentially suitable for specific P2Y1 receptor isolation

AU Zundorf, G.; Schafer, R.; Vohringer, C.; Halbfinger, E.; Fischer, B.; Reiser, G.

CS Medizinische Fakultät, Institut für Neurobiochemie, Otto-von-Guericke-Universität, Magdeburg, D-39120, Germany

SO Biochemical Pharmacology (2001), 61(10), 1259-1269

CODEN: BCPA6; ISSN: 0006-2952

PB Elsevier Science Inc.

DT Journal

LA English

AB Rat brain P2Y1 (rP2Y1) receptor-transfected human embryonic kidney cells (HEK 293) were recently shown to have enhanced reactivity to both ATP and ADP. Here, the authors demonstrated the usefulness of this cell line as a system for further studying novel adenine nucleotide analogs and for the biochem. characterization of the P2Y1 receptor. By measurement of intracellular Ca²⁺ release, for 2-butylthio-, 2-butylamino-, and 2-butyloxy-ATP (2-BuS-, 2-BuNH-, 2-BuO-ATP), EC₅₀ values of 1.3, 5, and 60 nM were determined, markedly lower than the value for ATP (130 nM). The EC₅₀ for 2-BuSADP was 1.1 nM. The corresponding 8-substituted ATP analogs showed a substantially lower potency than ATP (ATP > 8-BuSATP > 8-BuNHATP ≈ 8-BuOATP). AMP induced intracellular Ca²⁺ release with a very low potency; 2- and 8-substitutions on AMP caused no significant potency shift, except for 2-BuSAMP (EC₅₀ = 180 nM). Another new P2Y receptor probe, 2-[(6-biotinylamido)-hexylthio]ATP, was 22-fold more potent than ATP (EC₅₀ = 6 nM), revealing that even more bulky substituents linked to the C-2 position bind with high affinity at the P2Y1 receptor. This biotinylated probe was successfully used for the enrichment of the P2Y1 receptor tagged with green fluorescent protein from a crude membrane fraction. This one-step enrichment provides a substantial advance for P2Y1 receptor purification. Thus, human embryonic kidney 293 cells stably transfected with the rP2Y1 receptor represent a powerful model system for pharmacol. characterization of the P2Y1 receptor, circumventing problems associated with natural systems. They provide a means for the development of P2Y1 ligands of high potency and a good source for obtaining purified P2Y1 receptor.


IT 284040-54-6P 344402-39-7P

RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); PROC (Process)
(ATP analogs pharmacol. characterized in HEK293 cells expressing rat brain P2Y1 receptor in relation to biotinylated analog potentially suitable for specific P2Y1 receptor isolation)

RN 284040-54-6 CAPLUS


CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 344402-39-7 CAPLUS
 CN 5'-Adenylic acid, 8-(butylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 8 OF 17 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2000:304989 CAPLUS
 DN 133:105244
 TI Novel Inhibitors of Nucleoside Triphosphate Diphosphohydrolases: Chemical
 Synthesis and Biochemical and Pharmacological Characterizations
 AU Gendron, Fernand-Pierre; Halbfinger, Efrat; Fischer, Bilha; Duval,
 Martine; D'Orleans-Juste, Pedro; Beaudoin, Adrien R.
 CS Department de Biologie, Universite de Sherbrooke, Sherbrooke, QC, J1K 2R1,
 Can.
 SO Journal of Medicinal Chemistry (2000), 43(11), 2239-2247
 CODEN: JMCMAR; ISSN: 0022-2623
 PB American Chemical Society
 DT Journal
 LA English
 AB To elucidate the physiolo. role played by nucleoside triphosphate
 diphosphohydrolase (NTPDase; EC 3.6.1.5), adenine nucleotide analogs,
 modified on the purine ring, have been synthesized and tested as potential
 inhibitors. Resistance of ATP analogs to hydrolysis and their potency as
 NTPDase inhibitors were evaluated. For this purpose, a particulate
 fraction isolated from bovine spleen was used as the enzyme source. Among
 the synthesized analogs, 8-thiobutyladenosine 5'-triphosphate (8-BuS-ATP)
 was found to be the most effective nonhydrolyzable competitive inhibitor,
 with an estimated K_i of 10 μ M. This nonhydrolyzable analog did not exert
 any P2X-receptor-mediated effect on endothelium-denuded blood vessels,
 from the guinea pig mesenteric bed. In agreement with this observation,
 infusion of the analog did not cause any significant blood pressure
 variations of the precontracted vessel. Because in previous studies on
 isolated turkey erythrocytes and rat astrocytes 8-BuS-ATP was not able to
 trigger any P2Y1-receptor-mediated effect, it therefore appears that this
 NTPDase inhibitor does not interfere with purinergic receptors.
 IT 284040-53-5 284040-54-6
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological
 study, unclassified); BIOL (Biological study)
 (synthesis and biochem. and pharmacol. characterizations of novel


Printed

inhibitors of nucleoside triphosphate diphosphohydrolases)

RN 284040-53-5 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(hexylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-54-6 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 81609-35-0P 284040-51-3P 284040-52-4P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(synthesis and biochem. and pharmacol. characterizations of novel inhibitors of nucleoside triphosphate diphosphohydrolases)

RN 81609-35-0 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(ethylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-51-3 CAPLUS


CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(cycloheptylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 284040-52-4 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-[(2,2-dimethylpropyl)thio]-
 (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 61 THERE ARE 61 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 9 OF 17 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1999:771170 CAPLUS

DN 132:102410

TI Molecular Recognition of Modified Adenine Nucleotides by the
 P2Y1-Receptor. 1. A Synthetic, Biochemical, and NMR Approach

AU Halbfinger, Efrat; Major, Dan T.; Ritzmann, Marco; Ubl, Joachim; Reiser,
 Georg; Boyer, Jose L.; Harden, Kendall T.; Fischer, Bilha

CS Department of Chemistry Gonda-Goldschmied Center, Bar-Ilan University,
 Ramat-Gan, 52900, Israel

SO Journal of Medicinal Chemistry (1999), 42(26), 5325-5337
 CODEN: JMCMAR; ISSN: 0022-2623

PB American Chemical Society

DT Journal

LA English

AB The remarkably high potencies of 2-thioether-adenine nucleotides regarding
 the activation of the P2Y1-receptor (P2Y1-R) in turkey erythrocyte
 membranes represent some of the largest substitution-promoted increases in
 potencies over that of a natural receptor ligand. This paper describes
 the investigation regarding the origin of the high potency of these P2Y1-R
 ligands over that of ATP. For this study, an integrated approach was
 employed combining the synthesis of new ATP analogs, their biochem.
 evaluation, and their SAR anal. involving NMR expts. and theor. calcns.
 These expts. and calcns. were performed to elucidate the conformation and
 to evaluate the electronic nature of the investigated P2Y1-R ligands. ATP
 analogs synthesized included derivs. where C2 or C8 positions were
 substituted with electron-donating groups such as ethers, thioethers, or
 amines. The compds. were tested for their potency to induce
 P2Y1-R-mediated activation of phospholipase C in turkey erythrocytes and
 Ca2+ response in rat astrocytes. 8-Substituted ATP and AMP derivs. had

little or no effect on phospholipase C or on calcium levels, whereas the corresponding 2-substituted ATP analogs potentially increased the levels of inositol phosphates and $[Ca^{2+}]_i$. AMP analogs were ineffective except for 2-butylthio-AMP which induced a small Ca^{2+} response. P2Y1-R activity of these compds. was demonstrated by testing these ligands also on NG108-15 neuroblastoma + glioma hybrid cells. NMR data together with theor. calcns. imply that steric, rather than electronic, effects play a major role in ligand binding to the P2Y1-R. Hydrophobic interactions and H-bonds of the C2 substituent appear to be important determinants of a P2Y1-R ligand affinity.

IT 71683-16-4P 255716-10-0P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(preparation and structure activity relations of modified adenine nucleotides as P2Y1 receptor agonists)

RN 71683-16-4 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)-, tetrasodium salt (9CI) (CA INDEX NAME)

Absolute stereochemistry.




● 4 Na

RN 255716-10-0 CAPLUS

CN 5'-Adenylic acid, 8-(butylthio)-, disodium salt (9CI) (CA INDEX NAME)

Absolute stereochemistry.



● 2 Na

RE.CNT 51 THERE ARE 51 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 10 OF 17 CAPLUS COPYRIGHT 2006 ACS on STN


AN 1997:804940 CAPLUS

DN 128:164073

TI Quantitative one step derivatization of oligonucleotides by a fluorescent label through abasic site formation

AU Boturyn, Didier; Defrancq, Eric; Ducros, Veronique; Fontaine, Catherine; Lhomme, Jean
 CS LEDSS, UMR 5616, Universite Joseph Fourier - Grenoble I, GRENOBLE, 38041, Fr.
 SO Nucleosides & Nucleotides (1997), 16(10 & 11), 2069-2077
 CODEN: NUNUD5; ISSN: 0732-8311
 PB Marcel Dekker, Inc.
 DT Journal
 LA English
 AB Reaction of abasic site-containing oligonucleotides with an oxyamino fluorescent label is described. The reaction represents an efficient method to functionalize oligonucleotides at preselected positions.
 IT 157999-80-9
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (quant. one step derivatization of oligonucleotides by a fluorescent label through abasic site formation)
 RN 157999-80-9 CAPLUS
 CN Adenosine, 2'-deoxyguanylyl-(3'→5')-2'-deoxy-8-(propylthio)adenylyl-(3'→5')-2'-deoxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.




IT 202598-24-1P 202598-26-3P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (quant. one step derivatization of oligonucleotides by a fluorescent label through abasic site formation)
 RN 202598-24-1 CAPLUS
 CN Adenosine, 2'-deoxyguanylyl-(3'→5')-2'-deoxy-8-[(R)-propylsulfinyl]adenylyl-(3'→5')-2'-deoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A




PAGE 1-B




RN 202598-26-3 CAPLUS
CN Adenosine, 2'-deoxyguanylyl-(3'→5')-2'-deoxy-8-[(S)-propylsulfinyl]adenylyl-(3'→5')-2'-deoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A







RE.CNT 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 11 OF 17 CAPLUS COPYRIGHT 2006 ACS on STM
AN 1994:605872 CAPLUS
DN 121:205872
TI Hydrolysis of oligodeoxyribonucleotides containing 8-substituted purine nucleosides. A new route for preparing abasic oligodeoxyribonucleotides
AU Laayoun, Ali; Decout, Jean-Luc; Defrancq, Eric; Lhomme, Jean
CS L.E.D.S.S., URA CNRS, Univ. Joseph Fourier, Grenoble, 38041, Fr.
SO Tetrahedron Letters (1994), 35(28), 4991-4
CODEN: TELEAY; ISSN: 0040-4039
DT Journal
LA English
AB 2'-Deoxyadenosine substituted at C-8 by a propylthio group was introduced into oligodeoxyribonucleotides by solid phase synthesis. Oxidation by potassium persulfate occurred selectively on the sulfur containing nucleoside causing a weakening of the glycosidic bond. Subsequent hydrolytic treatment led to selective removal of the modified base and generation of an abasic site. This constitutes a novel and convenient route for the chemical synthesis of oligodeoxyribonucleotides containing an abasic site at a preselected position in the sequence.
IT 157999-80-9P 158020-58-7P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and oxidation of)
RN 157999-80-9 CAPLUS
CN Adenosine, 2'-deoxyguanylyl-(3'→5')-2'-deoxy-8-(propylthio)adenylyl-(3'→5')-2'-deoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.









RN 158020-58-7 CAPLUS

CN Thymidine, 2'-deoxyadenylyl-(3'→5')-2'-deoxyguanylyl-(3'→5')-
2'-deoxycytidylyl-(3'→5')-2'-deoxy-8-(propylthio)adenylyl-
(3'→5')-2'-deoxyguanylyl-(3'→5')-2'-deoxyadenylyl-
(3'→5')- (9CI) (CA INDEX NAME)

Absolute stereochemistry.





L8 ANSWER 12 OF 17 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1982:616611 CAPLUS

DN 97:216611

TI Species- or isozyme-selective enzyme inhibitors. 8. Synthesis of disubstituted two-substrate condensation products as inhibitors of rat adenylate kinases

AU Kappler, Francis; Hai, Ton T.; Abo, Masanobu; Hampton, Alexander

CS Inst. Cancer Res., Fox Chase Cancer Cent., Philadelphia, PA, 19111, USA

SO Journal of Medicinal Chemistry (1982), 25(10), 1179-84

reads on parent case

CODEN: JMCMAR; ISSN: 0022-2623

DT Journal
LA English

AB Syntheses are described of 5'(R)- and 5'(S)-C-Me-ATP, 5'(R)- and 5'(S)-C-n-Pr-ATP, and the phosphonate isostere of ATP with a C(5')-CH₂-P system. Two of the five compds. inhibited rat muscle adenylate kinase (AK-M) 8-9.5 times more effectively than AK II (present in poorly differentiated rat hepatoma tissue) and the two other compds. inhibited AK II at least 2-fold more effectively than AK-M. P1-[8-Ethylthio adenosine-5']-P5-(adenosine-5') pentaphosphate (8-SEt-Ap5A) is a potent dual substrate site inhibitor of the rat isozymes with selectivity for AK II. Three derivs. of 8-SEt-Ap5A were synthesized: p1-[8-(ethylthio) adenosine-5']-p5-[5'(R)-C-methyladenosine-5'] pentaphosphate (I), its 5'(R)-C-n-Pr analog (II), and di(8-SEt)-Ap5A (III). I and II, are readily accessible via reaction of a derivative of ATP γ -piperidinate with an ADP derivative. Except in the interaction of III with AK-M, I-III acted as two-site competitive inhibitors of AK-M and AK II. Inhibitory potencies of I-III with the two isozymes varied over more than a 95-fold range, and inhibitory potencies for AK-M relative to those of AK II varied more than 61-fold. III was an effective inhibitor of AK II and exhibited at least 4 times more selectivity for AK II than 8-SEt-Ap5A.

IT 83683-78-7

RL: RCT (Reactant); RACT (Reactant or reagent)
(condensation reaction of, with ADP derivative)

RN 83683-78-7 CAPLUS


CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(ethylthio)-, compd. with N,N-dibutyl-1-butanamine (1:4) (9CI) (CA INDEX NAME)

CM 1

CRN 81609-35-0

CMF C12 H20 N5 O13 P3 S


Absolute stereochemistry.



CM 2

CRN 102-82-9

CMF C12 H27 N




IT 83683-81-2P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and condensation reaction of, with ADP derivative)

RN 83683-81-2 CAPLUS


CN Adenosine 5'-(trihydrogen diphosphate), 8-(ethylthio)-, P'-anhydride with 1-piperidinylphosphonic acid (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 83683-76-5P 83683-77-6P 83694-37-5P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation and inhibition of adenylate kinases by)
 RN 83683-76-5 CAPLUS
 CN Adenosine 5'-(hexahydrogen pentaphosphate), 8-(ethylthio)-,
 P''''→5'-ester with 9-(6-deoxy-β-D-allofuranosyl)-9H-purin-6-
 amine, pentasodium salt (9CI) (CA INDEX NAME)

PAGE 1-A





● 5 Na

PAGE 1-B




RN 83683-77-6 CAPLUS
 CN Adenosine 5'-(hexahydrogen pentaphosphate), 8-(ethylthio)-,
 P''''→5' ester with 8-(ethylthio)adenosine (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 83694-37-5 CAPLUS
 CN Adenosine 5'-(hexahydrogen pentaphosphate), 8-(ethylthio)-,
 P''''->5'-ester with 9-(6,7,8-trideoxy- β -D-allo-octofuranosyl)-
 9H-purin-6-amine, pentasodium salt (9CI) (CA INDEX NAME)




● 5 Na



L8 ANSWER 13 OF 17 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 1982:611265 CAPLUS
 DN 97:211265
 TI Species- or isozyme-specific enzyme inhibitors. 9. Selective effects in inhibitions of rat pyruvate kinase isozymes by adenosine 5'-diphosphate derivatives
 AU Hai, Ton T.; Abo, Masanobu; Hampton, Alexander
 CS Fox Chase Cancer Cent., Inst. Cancer Res., Philadelphia, PA, 19111, USA
 SO Journal of Medicinal Chemistry (1982), 25(10), 1184-8
 CODEN: JMCMAR; ISSN: 0022-2623
 DT Journal
 LA English
 AB Derivs. of ADP with a substituent of 1-4 atoms at any of 8 positions were synthesized and evaluated as substrates and inhibitors of the liver (L), muscle (M), and kidney (K) isoenzymes of rat pyruvate kinase (I). The inhibitory potencies of the compds. were expressed as $K_m(\text{ADP})/K_i$ or as $K_m(\text{ADP})/K_m$ when no K_i value was available. Nine of 14 ADP derivs. exhibited differential inhibitions. The M and K isoenzymes, which cross-react immunol. with each other, but not with the L form, were inhibited differentially by 5 of the 14 derivs. I-K was preferentially inhibited by 2 derivs., I-L by 3 derivs., and I-M by 2 derivs. Among the most selective and/or effective inhibitors were 3'-OMe-ADP [$K_m(\text{ADP})/K_i = 0.07$ with I-K; inhibitory potency, K:M:L, 7.6:6.0:1], N6-Me,N6-(CH₂)₄N(Me)COMe-ADP (prepared previously) [$K_m(\text{ADP})/K_m = 0.43$ with I-L; inhibitory potency, L:K:M, 3:2:1], and 8-NHET-ADP [$K_m(\text{ADP})/K_i = 1.0$ with I-M; inhibitory potency, M:K:L, 7.1:1.2:1]. These and previous studies with 2 other enzymes indicate that monosubstituted substrate derivs. that bear short substituents (usually 1-4 atoms) at various positions are potentially useful probes in early stages of the attempted design of isoenzyme-selective inhibitors.
 IT **81609-45-2P**
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation and reaction kinetics with pyruvate kinase isoenzymes)
 RN 81609-45-2 CAPLUS
 CN Adenosine 5'-(trihydrogen diphosphate), 8-(ethylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L8 ANSWER 14 OF 17 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 1982:402705 CAPLUS
 DN 97:2705
 TI Species- or isozyme-specific enzyme inhibitors. 7. Selective effects in inhibitions of rat adenylate kinase isozymes by adenosine 5'-phosphate derivatives
 AU Hai, Ton T.; Picker, Donald; Abo, Masanobu; Hampton, Alexander
 CS Fox Chase Cancer Cent., Inst. Cancer Res., Philadelphia, PA, 19111, USA
 SO Journal of Medicinal Chemistry (1982), 25(7), 806-12
 CODEN: JMCMAR; ISSN: 0022-2623
 DT Journal
 LA English
 AB Monosubstituted derivs. of AMP with substituents of 1-3 atoms or group replacements at any of 11 positions were synthesized and examined as substrates and inhibitors of the rat muscle adenylate kinase isoenzyme

(AK-M) and the rat AK II and III isoenzymes predominant in poorly differentiated hepatoma tissue and normal liver tissue, resp. Inhibition indexes of the compds. were expressed as $K_m(\text{AMP})/K_i$ for competitive inhibition or as $K_m(\text{AMP})/K_m$ when only K_m was available. Substituents at N(1), N(6), or C(8) or on the ionizable phosphate O atom reduced inhibition below measurable levels; 2'-deoxy-AMP and adenosine 5'-sulfate had identical inhibition indexes with all 3 isoenzymes; compds. with substituents at C(2), O(2'), O(3'), C(4'), C(5'), or O(5') had higher inhibition indexes with AK-M than with AK II or III, and the same or similar indexes for AK II and III. The most effective and(or) selective inhibitors were 2-NHMe-AMP (index with AK-M, 0.2; index ratio, AK-M/AK III, 9.1), 2'-O-Me-AMP (index with AK-M, 0.14; index ratio, AK-M/AK III, 8.2), 2',3'-O-CMe₂-AMP (index with AK-M, 0.25; index ratio, AK-M/AK II, 6.6), 4'-allyl-AMP (index with AK-M, 0.97; index ratio, AK-M/AK III, 8.1), and 5'(S)-Et-AMP (index with AK-M, 0.64; index ratio, AK-M/AK II, 11.2). The study provided addnl. evidence that the attachment of simple substituents to various atoms in turn of a substrate is a potentially useful approach in early stages of the attempted design of isoenzyme-selective inhibitors.

IT 81921-37-1


RL: BIOL (Biological study)

(adenylate kinase isoenzyme inhibition by, structure in relation to)

RN 81921-37-1 CAPLUS

CN 5'-Adenylic acid, 8-(ethylthio)-, disodium salt (9CI) (CA INDEX NAME)

Absolute stereochemistry.



● 2 Na

L8 ANSWER 15 OF 17 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1982:402685 CAPLUS

DN 97:2685

TI Species- or isozyme-specific enzyme inhibitors. 4. Design of a two-site inhibitor of adenylate kinase with isozyme selectivity

AU Hampton, Alexander; Kappler, Francis; Picker, Donald

CS Inst. Cancer Res., Fox Chase Cancer Cent., Philadelphia, PA, 19111, USA

SO Journal of Medicinal Chemistry (1982), 25(6), 638-44

~~CODEN: JMCMAR; ISSN: 0022-2623~~

DT Journal

LA English


AB The ATP analogs, 6-butylamino-, 6-dibutylamino-, and 6-butylthio-9 β -D-ribofuranosylpurine 5'-triphosphate (I, II, and III, resp.), were synthesized and studied as inhibitors and(or) substrates of the rat muscle adenylate kinase isoenzyme (AK M) and the rat liver isoenzymes AK II and III. I and III were substrates (V_{max} relative to ATP, 13-190%) of the 3 AK isoenzymes, whereas II was a weak substrate and a competitive inhibitor of AK M and AK III. The affinities of the analogs relative to ATP [$K_m(\text{ATP})/K_m$ or K_i] were 0.03-0.075 for AK III and 0.14-0.28 for AK M, and the affinities for AK M exceeded those for AK III by factors of 2.3-7.0. Ap5A was synthesized by an improved method and was found to be a potent 2-site inhibitor ($K_i = 0.28 \mu\text{M}$), competitive toward AMP or ATP, for the 3 AK isoenzymes. 8-Ethylthio-Ap5A (IV) also behaved as a 2-site inhibitor; the 8-ethylthio group reduced the affinity for AK M 12-fold, but increased the

affinity for AK II and III 4-fold, resulting in .apprx.45-fold more effective inhibition of AK II and III ($K_i = 0.07 \mu\text{M}$) than of AK M ($K_i = 3.25 \mu\text{M}$). The 8-ethylthio group of 8-ethylthio-ATP (V) likewise reduced the affinity for the ATP site of AK M, but enhanced the affinity for the ATP sites of AK II and III, resulting in ≥ 30 -fold more effective inhibition of AK II and III. 8-Ethylthio-AMP inhibited AK II and III noncompetitively ($K_i = 21\text{-}24 \text{ mM}$) with respect to AMP, indicating that the 8-ethylthioadenosine moiety of IV probably binds to the ATP sites of these isoenzymes. IV had .apprx.1000-fold more affinity for AK II or III than did V. The findings indicate that isoenzyme-selective inhibitory effects of a substrate derivative can be imparted to a 2-site inhibitor, leading to significant enhancement of inhibitory potency.

IT 81609-34-9P 81609-35-0P 81609-36-1P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation and adenylate kinase isoenzyme inhibition by)
 RN 81609-34-9 CAPLUS
 CN Adenosine 5'-(hexahydrogen pentaphosphate), 8-(ethylthio)-,
 P'''' \rightarrow 5'-ester with adenosine, pentasodium salt (9CI) (CA INDEX
 NAME)


Absolute stereochemistry.

PAGE 1-A




● 5 Na

PAGE 1-B




RN 81609-35-0 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(ethylthio)- (9CI) (CA INDEX
 NAME)

Absolute stereochemistry.



RN 81609-36-1 CAPLUS
 CN 5'-Adenylic acid, 8-(ethylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 81609-46-3
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (reaction of, with tributylammonium adenosine trimetaphosphate)
 RN 81609-46-3 CAPLUS
 CN Adenosine 5'-(trihydrogen diphosphate), 8-(ethylthio)-, compd. with
 N,N-dibutyl-1-butanamine (1:1) (9CI) (CA INDEX NAME)

CM 1


CRN 81609-45-2
 CMF C12 H19 N5 O10 P2 S

Absolute stereochemistry.




CM 2


CRN 102-82-9
 CMF C12 H27 N



L8 ANSWER 16 OF 17 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 1981:525864 CAPLUS
 DN 95:125864
 TI Inhibition of inosinic acid dehydrogenase by 8-substituted purine nucleotides
 AU Skibo, Edward B.; Meyer, Rich B., Jr.
 CS Sch. Pharm., Univ. California, San Francisco, CA, 94143, USA
 SO Journal of Medicinal Chemistry (1981), 24(10), 1155-61
 CODEN: JMCMAR; ISSN: 0022-2623
 DT Journal
 LA English
 GI



I



II


AB Twenty-seven AMP and IMP derivs. I and II (R = CH₂Ph, substituted benzyl, CH₂CH₂Ph, etc.) were synthesized and studied for Escherichia coli IMP dehydrogenase (EC 1.2.1.14) [9028-93-7] inhibiting activity. Many inhibitors of this enzyme have anticancer activity. All of the compds. studied were competitive inhibitors in IMP-dependent competition studies and lacked substrate activity. Multiple regression anal. showed that for I and II (R = para substituted benzyl), the electron-withdrawing ability of the substituent on the benzylthio moiety correlated best with the K_i of the analogs.

IT 78710-82-4P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation and IMP dehydrogenase inhibition by, antitumor activity in relation to)


RN 78710-82-4 CAPLUS

CN 5'-Adenylic acid, 8-(pentylthio)-, dilithium salt (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L8 ANSWER 17 OF 17 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 1979:604203 CAPLUS
 DN 91:204203
 TI Design of species- or isozyme-specific enzyme inhibitors. 3. Species and isozymic differences between mammalian and bacterial adenylate kinases in substituent tolerance in an enzyme-substrate complex
 AU Hampton, Alexander; Picker, Donald
 CS Fox Chase Cancer Cent., Inst. Cancer Res., Philadelphia, PA, 19111, USA
 SO Journal of Medicinal Chemistry (1979), 22(12), 1529-32
 CODEN: JMCMAR; ISSN: 0022-2623
 DT Journal
 LA English
 GI



AB The ATP derivs. I (R = alkylthio, hydroxyalkylthio, or PhS) were prepared from tetra-Li 8-bromoadenosine 5'-triphosphate [71683-13-1] and the appropriate mercaptide and converted to the tetra Na salts. I and N6-ATP derivs. were evaluated as potential species- or isoenzyme-selective inhibitors of bacterial and mammalian adenylate kinase [9013-02-9]. The substituent attached at either N6 or C-8 influenced the affinity of the compds. for the enzymic ATP sites in both a species- and an isoenzyme-selective manner. Structure-activity relations are discussed.


IT 71683-14-2P 71683-15-3P 71683-16-4P
 71683-17-5P

RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation and adenylate kinase-inhibiting activity of)

RN 71683-14-2 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(ethylthio)-, tetrasodium salt (9CI) (CA INDEX NAME)

Absolute stereochemistry.




● 4 Na

RN 71683-15-3 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(propylthio)-, tetrasodium

salt (9CI) (CA INDEX NAME)

Absolute stereochemistry.




●4 Na

RN 71683-16-4 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)-, tetrasodium salt (9CI) (CA INDEX NAME)

Absolute stereochemistry.




●4 Na

RN 71683-17-5 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(pentylthio)-, tetrasodium salt (9CI) (CA INDEX NAME)

Absolute stereochemistry.



●4 Na

Connecting via Winsock to STN

10/620,520

Welcome to STN International! Enter x:x

LOGINID:sssptal600txm

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

- NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 JAN 17 Pre-1988 INPI data added to MARPAT
NEWS 4 FEB 21 STN AnaVist, Version 1.1, lets you share your STN AnaVist
visualization results
NEWS 5 FEB 22 The IPC thesaurus added to additional patent databases on STN
NEWS 6 FEB 22 Updates in EPFULL; IPC 8 enhancements added
NEWS 7 FEB 27 New STN AnaVist pricing effective March 1, 2006
NEWS 8 MAR 03 Updates in PATDPA; addition of IPC 8 data without attributes
NEWS 9 MAR 22 EMBASE is now updated on a daily basis
NEWS 10 APR 03 New IPC 8 fields and IPC thesaurus added to PATDPAFULL
NEWS 11 APR 03 Bibliographic data updates resume; new IPC 8 fields and IPC
thesaurus added in PCTFULL
NEWS 12 APR 04 STN AnaVist \$500 visualization usage credit offered
NEWS 13 APR 12 LINSPEC, learning database for INSPEC, reloaded and enhanced
NEWS 14 APR 12 Improved structure highlighting in FQHIT and QHIT display
in MARPAT
NEWS 15 APR 12 Derwent World Patents Index to be reloaded and enhanced during
second quarter; strategies may be affected
NEWS 16 MAY 10 CA/CAPplus enhanced with 1900-1906 U.S. patent records
NEWS 17 MAY 11 KOREAPAT updates resume
NEWS 18 MAY 19 Derwent World Patents Index to be reloaded and enhanced
- NEWS EXPRESS FEBRUARY 15 CURRENT VERSION FOR WINDOWS IS V8.01a,
CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 19 DECEMBER 2005.
V8.0 AND V8.01 USERS CAN OBTAIN THE UPGRADE TO V8.01a AT
<http://download.cas.org/express/v8.0-Discover/>
- NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items
NEWS IPC8 For general information regarding STN implementation of IPC 8
NEWS X25 X.25 communication option no longer available after June 2006

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

* * * * *

COMPLETE THE STN SURVEY - APRIL 27 THROUGH MAY 31

Dear valued STN customer,

In an effort to enhance your experience with STN, we would
like to better understand what you find useful. Please take
approximately 5 minutes to complete a web survey.

If you provide us with your name, login ID, and e-mail address, you will be entered in a drawing to win a free iPod(R). Your responses will be kept confidential and will help us make future improvements to STN.

Take survey: <http://www.zoomerang.com/survey.zgi?p=WEB2259HNKWTUW>

Thank you in advance for your participation.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 12:31:35 ON 26 MAY 2006

=> file reg

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'REGISTRY' ENTERED AT 12:31:43 ON 26 MAY 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 25 MAY 2006 HIGHEST RN 885654-58-0

DICTIONARY FILE UPDATES: 25 MAY 2006 HIGHEST RN 885654-58-0

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 6, 2006

Please note that search-term pricing does apply when conducting SmartSELECT searches.

*
* The CA roles and document type information have been removed from *
* the IDE default display format and the ED field has been added, *
* effective March 20, 2005. A new display format, IDERL, is now *
* available and contains the CA role and document type information. *
*

Structure search iteration limits have been increased. See HELP SLIMITS for details.


REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

=>
Uploading C:\Program Files\Stnexp\Queries\10620520.str

L1 STRUCTURE UPLOADED

=> d l1
L1 HAS NO ANSWERS
L1 STR



G1 Ak

Structure attributes must be viewed using STN Express query preparation.

=> s l1 sam

SAMPLE SEARCH INITIATED 12:32:03 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 3555 TO ITERATE

56.3% PROCESSED 2000 ITERATIONS 50 ANSWERS
INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 67525 TO 74675
PROJECTED ANSWERS: 1272 TO 2424

L2 50 SEA SSS SAM L1

=> s l1 full

FULL SEARCH INITIATED 12:32:11 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 73256 TO ITERATE

100.0% PROCESSED 73256 ITERATIONS 1324 ANSWERS
SEARCH TIME: 00.00.01

L3 1324 SEA SSS FUL L1

=> file caplus

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	166.94	167.15

FILE 'CAPLUS' ENTERED AT 12:32:17 ON 26 MAY 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is

strictly prohibited.

FILE COVERS 1907 - 26 May 2006 VOL 144 ISS 23
FILE LAST UPDATED: 25 May 2006 (20060525/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply.
They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> s l3

L4 1225 L3

=> s ntpdase

114 NTPDASE
48 NTPDASES
L5 130 NTPDASE
(NTPDASE OR NTPDASES)

=> s nucleoside triphosphate diphosphohydrolase

45789 NUCLEOSIDE
30934 NUCLEOSIDES
57166 NUCLEOSIDE
(NUCLEOSIDE OR NUCLEOSIDES)
39515 TRIPHOSPHATE
10190 TRIPHOSPHATES
45662 TRIPHOSPHATE
(TRIPHOSPHATE OR TRIPHOSPHATES)
442 DIPHOSPHOHYDROLASE
50 DIPHOSPHOHYDROLASES
452 DIPHOSPHOHYDROLASE
(DIPHOSPHOHYDROLASE OR DIPHOSPHOHYDROLASES)
L6 115 NUCLEOSIDE TRIPHOSPHATE DIPHOSPHOHYDROLASE
(NUCLEOSIDE (W) TRIPHOSPHATE (W) DIPHOSPHOHYDROLASE)

=> s ectonucleotidase

293 ECTONUCLEOTIDASE
157 ECTONUCLEOTIDASES
L7 389 ECTONUCLEOTIDASE
(ECTONUCLEOTIDASE OR ECTONUCLEOTIDASES)

=> s atpdase

58 ATPDASE
15 ATPDASES
L8 60 ATPDASE
(ATPDASE OR ATPDASES)

=> s e-type atpase

1936381 E
1677630 TYPE
580656 TYPES
2126695 TYPE
(TYPE OR TYPES)
81743 ATPASE
7090 ATPASES
82810 ATPASE
(ATPASE OR ATPASES)
L9 20 E-TYPE ATPASE
(E (W) TYPE (W) ATPASE)

=> s ecto atpase

3357 ECTO
2 ECTOS
3359 ECTO
(ECTO OR ECTOS)
81743 ATPASE
7090 ATPASES
82810 ATPASE
(ATPASE OR ATPASES)

L10 457 ECTO ATPASE
(ECTO(W)ATPASE)

=> d his

(FILE 'HOME' ENTERED AT 12:31:35 ON 26 MAY 2006)

FILE 'REGISTRY' ENTERED AT 12:31:43 ON 26 MAY 2006

L1 STRUCTURE UPLOADED
L2 50 S L1 SAM
L3 1324 S L1 FULL

FILE 'CAPLUS' ENTERED AT 12:32:17 ON 26 MAY 2006

L4 1225 S L3
L5 130 S NTPDASE
L6 115 S NUCLEOSIDE TRIPHOSPHATE DIPHOSPHOHYDROLASE
L7 389 S ECTONUCLEOTIDASE
L8 60 S ATPDASE
L9 20 S E-TYPE ATPASE
L10 457 S ECTO ATPASE

=> s l4 and (l5 or l6 or l7 or l8 or l9 or l10)

L11 11 L4 AND (L5 OR L6 OR L7 OR L8 OR L9 OR L10)

=> d bib abs hitstr 1-11 l11

L11 ANSWER 1 OF 11 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2004:911398 CAPLUS

DN 142:214281

TI C8-substituted purine nucleotide analogs and their use as inhibitors of
nucleoside triphosphate diphosphohydrolases

IN Halbfinger, Efrat; Fischer, Bilha; Beaudoin, Adrien R.; Gendron, Fernand
Pierre

PA Universite de Sherbrooke, Can.; Bar-Ilan University

SO Can. Pat. Appl., 54 pp.

CODEN: CPXXEB

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	CA 2311084	AA	20011209	CA 2000-2311084	20000609
PRAI	CA 2000-2311084		20000609		

OS CASREACT 142:214281

AB Ectonucleoside triphosphate diphosphohydrolases [NTPDases; EC 3.6.1.5] constitute a family of enzymes which are involved in the metabolism of extracellular nucleotides, catalyzing the hydrolysis of the gamma and beta phosphate bonds of triphospho- and diphosphonucleosides (whereas 5'-nucleotidases [EC 3.1.3.5] catalyze the hydrolysis of alpha phosphate bond of monophosphonucleosides). These extracellular nucleotides interact with endothelial, epithelial and smooth muscle cells, as well as blood cells and lymphoid cells, to influence the different physiol. systems of vertebrates. Since these ecto-nucleotidases alter the extracellular concns. of nucleotides these enzymes modulate their physiol. effects, including, for example, platelet aggregation, heart function, control of vascular tone and inflammation reactions, electrolyte secretion and gastrointestinal motility, neurotransmission both in central and peripheral nervous systems, as well as other effects in other physiol. systems. This invention provides C8 substituted purine nucleotide analogs, such as ATP analogs, and further provides their use as inhibitors of NTPDases and thereby as tools to modulate the conversion of nucleotides into nucleoside derivs., and thus modulate the levels of these compds. Such modulation further provides for the modulation of the activity and function of many processes which are affected by these compds.

IT 81609-35-0P 284040-51-3P 284040-52-4P
284040-53-5P

RL: BSU (Biological study, unclassified); PAC (Pharmacological activity);
SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological

study); PREP (Preparation); USES (Uses)


(C8-substituted purine nucleotide analogs and their use as inhibitors of nucleoside triphosphate

diphosphohydrolases to modulate purine nucleotide levels and biol. processes)

RN 81609-35-0 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(ethylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-51-3 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(cycloheptylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-52-4 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-[(2,2-dimethylpropyl)thio]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-53-5 CAPLUS


CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(hexylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 284040-54-6 284040-60-4
RL: BSU (Biological study, unclassified); PAC (Pharmacological activity);
THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(C8-substituted purine nucleotide analogs and their use as inhibitors
of **nucleoside triphosphate**
diphosphohydrolases to modulate purine nucleotide levels and
biol. processes)
RN 284040-54-6 CAPLUS
CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX
NAME)

Absolute stereochemistry.




RN 284040-60-4 CAPLUS
CN Adenosine 5'-(tetrahydrogen triphosphate), 8-butoxy- (9CI) (CA INDEX
NAME)

Absolute stereochemistry.



IT 23567-97-7, 8-Bromo-ATP
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(substrate; C8-substituted purine nucleotide analogs and their use as
inhibitors of **nucleoside triphosphate**
diphosphohydrolases to modulate purine nucleotide levels and
biol. processes)
RN 23567-97-7 CAPLUS
CN Adenosine 5'-(tetrahydrogen triphosphate), 8-bromo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 2 OF 11 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2003:856092 CAPLUS

DN 139:333119

TI Ecto-nucleoside triphosphate

diphosphohydrolase inhibition-based methods for screening for a compound useful in the treatment or prevention of lymphocytic disorders, for inhibiting lymphocyte activity and preventing or treating lymphocytic disorders

IN Beaudoin, Adrien; Benrezzak, Ouhida

PA Bioflash Inc., Can.

SO PCT Int. Appl., 63 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003089664	A1	20031030	WO 2003-CA583	20030422
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	CA 2382768	AA	20031019	CA 2002-2382768	20020419
	CA 2479501	AA	20031030	CA 2003-2479501	20030422
	AU 2003226989	A1	20031103	AU 2003-226989	20030422
	US 2005164306	A1	20050728	US 2003-511133	20030422

PRAI CA 2002-2382768 A 20020419
WO 2003-CA583 W 20030422

AB The invention discloses a method of screening for a compound useful in the treatment of a disease or condition characterized by an immune cell disorder, wherein the cell expresses ecto-nucleoside triphosphate diphosphohydrolases (NTPDases), the method comprising contacting a candidate compound with NTPDase, wherein the candidate compound is selected if the activity of the NTPDase is reduced in the presence of the candidate compound as compared to that in the absence thereof. The invention also discloses a method for inhibiting an immune cell activity in a mammal, comprising targeting immune cells with an effective amount of a NTPDase inhibitor. The invention further discloses a method to prevent or reduce the risk of rejection of transplanted tissue or organ, comprising administering to the animal an effective amount of NTPDase inhibitor.

IT 284040-54-6 344402-39-7


RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(ecto-nucleoside triphosphate

**diphosphohydrolase inhibition-based methods for screening for
agents for treatment of immune cell disorder-associated conditions)**

RN 284040-54-6 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX
NAME)


Absolute stereochemistry.



RN 344402-39-7 CAPLUS

CN 5'-Adenylic acid, 8-(butylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 3 OF 11 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2003:711173 CAPLUS

DN 139:230955

TI Preparation of C8-substituted purine nucleotide analogs as **NTPDase**
inhibitors

IN Beaudoin, Adrien R.; Gendron, Fernand-Pierre; Halbfinger, Efrat; Fischer,
Bilha

PA Bar-Ilan University, Israel; Universite De Sherbrooke

SO U.S., 21 pp.

CODEN: USXXAM


DT Patent

LA English

FAN.CNT 1

Parent Case

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	US 6617439	B1	20030909	US 2000-591177	20000609
	US 2004043955	A1	20040304	US 2003-620520	20030716
PRAI	US 2000-591177	A3	20000609		
OS	MARPAT 139:230955				
GI					



AB C8-substituted purine nucleotide analogs, I (R is alkyl, cycloalkyl) such as ATP analogs, and their use is described, including their use as inhibitors of **NTPDases** and thereby as tools to modulate the conversion of nucleotides into nucleoside derivs., and thus modulate the levels of these compds. Such modulation further provides for the modulation of the activity and function of many processes which are affected by these compds. Thus, I [R = (CH₂)₃Me] was prepared and tested in vivo as NTPDase inhibitor. I [R = (CH₂)₃Me] interacts specifically with the binding site of the enzyme potentially reduces the risk of interference with other ATP-binding enzymes or receptors, and thus possesses a high degree of specificity. The compds. of the invention were analyzed with resp. to any effects on the activity of purinoceptors.


IT 23567-97-7, 8-Bromoadenosine triphosphate

RL: BSU (Biological study, unclassified); BIOL (Biological study)
(preparation of C8-substituted purine nucleotide analogs as **NTPDase** inhibitors)

RN 23567-97-7 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-bromo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 81609-35-0P 284040-51-3P 284040-52-4P

284040-53-5P 284040-54-6P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of C8-substituted purine nucleotide analogs as **NTPDase** inhibitors)

RN 81609-35-0 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(ethylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-51-3 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(cycloheptylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-52-4 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-[(2,2-dimethylpropyl)thio]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-53-5 CAPLUS


CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(hexylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 284040-54-6 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 284040-59-1 284040-60-4
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (preparation of C8-substituted purine nucleotide analogs as **NTPDase** inhibitors)
 RN 284040-59-1 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylamino)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 284040-60-4 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-butoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 71 THERE ARE 71 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 4 OF 11 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2002:910711 CAPLUS

DN 138:232038

TI Heteromultimerization modulates P2X receptor functions through participating extracellular and C-terminal subdomains

AU Koshimizu, Taka-aki; Ueno, Susumu; Tanoue, Akito; Yanagihara, Nobuyuki; Stojilkovic, Stanko S.; Tsujimoto, Gozoh

CS Department of Molecular and Cell Pharmacology, National Institutes of Health, NICHD, Bethesda, MD, 20892, USA

SO Journal of Biological Chemistry (2002) 277(49), 46891-46899

CODEN: JBCHA3; ISSN: 0021-9258

PB American Society for Biochemistry and Molecular Biology

DT Journal

LA English

AB P2X purinergic receptors (P2XRs) differ among themselves with respect to their ligand preferences and channel kinetics during activation, desensitization, and recovery. However, the contributions of distinct receptor subdomains to the subtype-specific behavior have been incompletely characterized. Homomeric receptors having the extracellular domain of the P2X3 subunit in the P2X2a-based backbone (P2X2a/X3ex) mimicked two intrinsic functions of P2X3R, sensitivity to $\alpha\beta$ -methylene ATP and **ecto-ATPase**-dependent recovery from endogenous desensitization; these two functions were localized to the N- and C-terminal halves of the P2X3 extracellular loop, resp. The chimeric P2X2aR/X3ex receptors also desensitized with accelerated rates compared with native P2X2aR, and the introduction of P2X2 C-terminal splicing into the chimeric subunit (P2X2b/X3ex) further increased the rate of desensitization. Phys. and functional heteromerization of native P2X2a and P2X2b subunits was also demonstrated. In heteromeric receptors, the ectodomain of P2X3 was a structural determinant for ligand selectivity and recovery from desensitization, and the C terminus of P2X2 was an important factor for the desensitization rate. Furthermore, [γ - 32 P]8-azido ATP, a photoreactive agonist, was effectively crosslinked to P2X3 subunit in homomeric receptors but not in heteromeric P2X2 + P2X3Rs. These results indicate that heteromeric receptors formed by distinct P2XR subunits develop new functions resulting from integrative effects of the participating extracellular and C-terminal subdomains.


IT 53696-59-6, 8-Azido ATP

RL: BSU (Biological study, unclassified); BIOL (Biological study)
(heteromultimerization modulates P2X receptor functions through participating extracellular and C-terminal subdomains as studied in GT1 cells)

RN 53696-59-6 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-azido- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 5 OF 11 CAPLUS COPYRIGHT 2006 ACS on STN
AN 2001:327844 CAPLUS
DN 135:149038
TI Inhibitors of **NTPDase**: key players in the metabolism of
extracellular purines
AU Gendron, F. P.; Halbfinger, E.; Fischer, B.; Beaudoin, A. R.
CS Department of Biology, University of Sherbrooke, Sherbrooke, Can.
SO Advances in Experimental Medicine and Biology (2000), 486(Purine and
Pyrimidine Metabolism in Man X), 119-123
CODEN: AEMBAP; ISSN: 0065-2598

PB Kluwer Academic/Plenum Publishers
DT Journal
LA English

AB This study described the potential of a new class of ATP analogs as
nucleoside triphosphate diphosphohydrolase (**NTPDase**) inhibitors. From previous studies, 8-thiobutyladenosine
5'-triphosphate (8-BuS-ATP) appears to be a specific and efficient
NTPDase inhibitor. This novel inhibitor is a new tool to regulate
NTPDase activity and thereby influencing purine signaling in
mammalian.


IT 23567-97-7 81609-35-0 284040-51-3
284040-53-5 284040-54-6 284040-59-1
284040-60-4 352690-39-2

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL
(Biological study); PROC (Process)
(inhibitors of **nucleoside triphosphate
diphosphohydrolase** - key players in metabolism of extracellular
purines)

RN 23567-97-7 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-bromo- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 81609-35-0 CAPLUS


CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(ethylthio)- (9CI) (CA INDEX
NAME)

Absolute stereochemistry.




RN 284040-51-3 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(cycloheptylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 284040-53-5 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(hexylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 284040-54-6 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-59-1 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylamino)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-60-4 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-butoxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 352690-39-2 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-[[[(1,1-dimethylethyl)thio]methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 6 OF 11 CAPLUS COPYRIGHT 2006 ACS on STN
AN 2000:304989 CAPLUS
DN 133:105244

TI Novel Inhibitors of **Nucleoside Triphosphate
Diphosphohydrolases**: Chemical Synthesis and Biochemical and
Pharmacological Characterizations

AU Gendron, Fernand-Pierre; Halbfinger, Efrat; Fischer, Bilha; Duval,
Martine; D'Orleans-Juste, Pedro; Beaudoin, Adrien R.

CS Department de Biologie, Universite de Sherbrooke, Sherbrooke, QC, J1K 2R1,
Can.

SO Journal of Medicinal Chemistry (2000), 43(11), 2239-2247

CODEN: JMCMAR; ISSN: 0022-2623

PB American Chemical Society

DT Journal

LA English

AB To elucidate the physiolo. role played by **nucleoside
triphosphate diphosphohydrolase (NTPDase; EC**

Printed

3.6.1.5), adenine nucleotide analogs, modified on the purine ring, have
been synthesized and tested as potential inhibitors. Resistance of ATP
analogs to hydrolysis and their potency as **NTPDase** inhibitors
were evaluated. For this purpose, a particulate fraction isolated from
bovine spleen was used as the enzyme source. Among the synthesized
analogs, 8-thiobutyladenosine 5'-triphosphate (8-BuS-ATP) was found to be
the most effective nonhydrolyzable competitive inhibitor, with an estimated K_i
of 10 μ M. This nonhydrolyzable analog did not exert any
P2X-receptor-mediated effect on endothelium-denuded blood vessels, from
the guinea pig mesenteric bed. In agreement with this observation,
infusion of the analog did not cause any significant blood pressure
variations of the precontracted vessel. Because in previous studies on
isolated turkey erythrocytes and rat astrocytes 8-BuS-ATP was not able to
trigger any P2Y1-receptor-mediated effect, it therefore appears that this
NTPDase inhibitor does not interfere with purinergic receptors.

IT 284040-53-5 284040-54-6 284040-59-1
284040-60-4


RL: BAC (Biological activity or effector, except adverse); BSU (Biological
study, unclassified); BIOL (Biological study)

(synthesis and biochem. and pharmacol. characterizations of novel
inhibitors of **nucleoside triphosphate
diphosphohydrolases**)

RN 284040-53-5 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(hexylthio)- (9CI) (CA INDEX
NAME)


Absolute stereochemistry.



RN 284040-54-6 CAPLUS


CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX
NAME)

Absolute stereochemistry.




RN 284040-59-1 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylamino)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 284040-60-4 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-butoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 81609-35-0P 284040-51-3P 284040-52-4P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (synthesis and biochem. and pharmacol. characterizations of novel inhibitors of nucleoside triphosphate diphosphohydrolases)
 RN 81609-35-0 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(ethylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 284040-51-3 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(cycloheptylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 284040-52-4 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-[(2,2-dimethylpropyl)thio]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 61 THERE ARE 61 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 7 OF 11 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1997:285894 CAPLUS

DN 126:340339

TI Inhibition of **ecto-ATPase** by the P2 purinoceptor agonists, ATP γ S, α,β -methylene-ATP, and AMP-PNP, in endothelial cells

AU Chen, Bing Chang; Lin, Wan-Wan

CS Department of Pharmacology, College of Medicine, National Taiwan University, Taipei, Taiwan

SO Biochemical and Biophysical Research Communications (1997), 233(2), 442-446

Printed

PB Academic
 DT Journal
 LA English
 AB

Ecto-ATPase is a plasma membrane-bound enzyme that sequentially dephosphorylates extracellular nucleotides such as ATP. This breakdown of ATP and other nucleotides makes it difficult to characterize and classify P2 purinoceptors. We have previously shown that the P2 purinergic antagonists, PPADS, suramin and reactive blue, act as **ecto-ATPase** inhibitors in various cell lines. Here, we show that the P2 purinergic agonists, ATPyS, α,β -methylene ATP (α,β -MeATP) and AMP-PNP, inhibit the **ecto-ATPase** of bovine pulmonary artery endothelial cells (CPAE), with pIC50 values of 5.2, 4.5 and 4.0, resp. In CPAE, FPL67156, a selective **ecto-ATPase** inhibitor, also inhibits **ecto-ATPase** activity, with a pIC50 value of 4.0. In addition, α,β -MeATP (3-100 μ M), which itself does not induce phosphoinositide (PI) turnover, left-shifted the agonist-concentration effect (E/[A]) curves for ATP, 2MeS-ATP and UTP by approx. 100-300 fold, while those for ATPyS and AMP-PNP were only shifted approx. 2-3 fold. Moreover, in the presence of α,β -MeATP, not only was the potentiation effect of PPADS on the UTP response lost, but a slight inhibition of the UTP response by PPADS was also seen. Thus, we conclude that the action of ATPyS, α,β -MeATP and AMP-PNP as **ecto-ATPase** inhibitors account for their high agonist potency, and also provide information for the development of **ecto-ATPase** inhibitors of high selectivity and potency.

IT 23567-97-7, 8-Bromo-ATP 53602-90-7, 8-(6-Aminoethyl)amino-ATP


RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(inhibition of **ecto-ATPase** by P2 purinoceptor agonists, ATPyS, α,β -methylene-ATP, and AMP-PNP, in endothelial cells)

RN 23567-97-7 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-bromo- (9CI). (CA INDEX NAME)


Absolute stereochemistry.



RN 53602-90-7 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-[(6-aminoethyl)amino]- (9CI)
 (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 8 OF 11 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1996:300566 CAPLUS

DN 124:336420

TI Hydrolysis of P2-purinoceptor agonists by a purified
ectonucleotidase from the bovine aorta, the ATP-diphosphohydrolase

AU Picher, Maryse; Sevigny, Jean; D'Orleans-Juste, Pedro; Beaudoin, Adrien R.

CS Fac. Sci., Univ. Sherbrooke, Sherbrooke, QC, Can.

SO Biochemical Pharmacology (1996), 51(11), 1453-1460

CODEN: BCPA6; ISSN: 0006-2952

PB Elsevier

DT Journal

LA English

AB Pharmacologists are becoming more and more aware of the possibility that certain ATP analogs currently used to classify the P2-purinoceptors are dephosphorylated by **ectonucleotidases**. In this study, the authors provide evidence that in the vascular system, these purine analogs are hydrolyzed by an ATP-diphosphohydrolase (**ATPDase**). This enzyme is known as the major plasma membrane nucleotidase of endothelial and smooth muscle cells, and it believed to dephosphorylate extracellular triphospho- and diphosphonucleosides. Assays were conducted with a purified **ATPDase** from smooth muscle cells of bovine aorta. At a concentration of 250 μM , adenosine 5'-(α,β -methylene) triphosphonate (α,β -metATP), adenosine 5'-(β,γ -methylene) triphosphonate (β,γ -metATP), adenosine 5'-(α,β -methylene) diphosphonate (α,β -metADP), adenylyl 5'-(β,γ -imido) diphosphonate (β,γ -metATP), adenosine 5'-O-(2-thiodiphosphate) (ADP βS) all resisted dephosphorylation, whereas 2-chloroadenosine triphosphate (2-chloroATP), 2-methylthioadenosine triphosphate (2-MeSATP) and 8-bromoadenosine triphosphate (8-bromoATP) were hydrolyzed at 99, 63, and 20% of the rate of ATP hydrolysis, resp. All the non-hydrolyzable analogs tested, except α,β -metADP, competed with ATP and ADP for the **ATPDase** catalytic site, reducing their hydrolysis by 35-50%. Apparent K_m values for ATP and ADP were estimated at 14.1 and 12.0 μM , resp., whereas apparent K_m and K_i values for the purine analogs ranged from 12 to 28 μM . These results strongly support the view that (1) the **ATPDase** is expected to reduce substantially the P2-response induced by ATP, ADP, and some hydrolyzable agonists; and (2) by competing with the hydrolysis of endogenously released ATP and ADP, non-hydrolyzable analogs could alter the amplitude or direction of the cellular response induced by these natural substrates.

IT 23567-97-7, 8-Bromoadenosine triphosphate


RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(hydrolysis of P2-purinoceptor agonists by ATP-diphosphohydrolase from bovine aorta)

RN 23567-97-7 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-bromo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 9 OF 11 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1994:50794 CAPLUS

DN 120:50794

TI 8-Azido-adenine nucleotides as substrates of ecto-nucleotidases in chromaffin cells: Inhibitory effect of photoactivation

AU Rodriguez-Pascual, Fernando; Torres, Magdalena; Miras-Portugal, M. Teresa

CS Fac. Vet., Univ. Complutense Madrid, Madrid, Spain

SO Archives of Biochemistry and Biophysics (1993), 306(2), 420-6

CODEN: ABBIA4; ISSN: 0003-9861

DT Journal

LA English

AB The components of the ecto-nucleotidase pathway at the extracellular surface of adrenal chromaffin cells are the enzymic activities responsible for the hydrolysis of granular nucleotide compds. released during the secretory response. The azido-nucleotides have been largely employed to characterize nucleotide binding sites. The 8-azido-adenine nucleotides were studied as substrates of ecto-nucleotidases in cultured chromaffin cells by HPLC procedures. 8-Azido-ATP (8-N3-ATP) was a good substrate for **ecto-ATPase** activity, the K_m value was $256.30 \pm 36.41 \mu M$, and the V_{max} value was $14.33 \pm 0.84 \text{ nmol/min} + 106 \text{ cells}$. 8-Azido-ADP (8-N3-ADP) was dephosphorylated by the ecto-ADPase activity with a K_m value of $595.29 \pm 67.44 \mu M$ and V_{max} value of $6.86 \pm 0.45 \text{ nmol/min} + 106 \text{ cells}$. These kinetic parameters were similar to those obtained with ATP and ADP in the same culture and incubation conditions. 8-Azido-AMP (8-N3-AMP) was not hydrolyzed by the ecto-5'-nucleotidase activity. The 8-azido-nucleotides competitively inhibited the hydrolysis of adenine nucleotides, with K_i values in the same range as the K_m . After UV photoactivation, the three 8-azido-nucleotides ($100 \mu M$) irreversibly inhibited and to a similar extent, between 40 and 55%, each of ecto-nucleotidase activities. UV photoactivation in the presence of nucleotides in the same concentration range was an effective protection from the inhibition.

IT 53696-59-6, 8-Azido-ATP 59432-65-4, 8-Azido-ADP

60731-47-7, 8-Azido-AMP


RL: BIOL (Biological study)

(ecto-nucleotidase specificity for, in adrenal medulla chromaffin cell, photoactivation study of)

RN 53696-59-6 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-azido- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



*already printed article w/
N-linked caps*


RN 59432-65-4 CAPLUS
CN Adenosine 5'-(trihydrogen diphosphate), 8-azido- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 60731-47-7 CAPLUS
CN 5'-Adenylic acid, 8-azido- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 10 OF 11 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1987:590370 CAPLUS

DN 107:190370

TI The structure-activity relationships of **ectonucleotidases** and of excitatory P2-purinoceptors: evidence that dephosphorylation of ATP analogs reduces pharmacological potency

AU Welford, Laurence A.; Cusack, Noel J.; Hourani, Susanna M. O.

CS King's Coll., Univ. London, London, WC2R 2LS, UK

SO European Journal of Pharmacology (1987), 141(1), 123-30

CODEN: EJPHAZ; ISSN: 0014-2999

DT Journal

LA English

AB The dephosphorylation of adenine nucleotides and their analogs by **ectonucleotidases** on the guinea pig urinary bladder was studied using HPLC. The rate of dephosphorylation of each analog was compared with its pharmacol. potency at causing contraction. ATP, ADP, and AMP were rapidly dephosphorylated, and substitution on the purine ring did not affect the rate of breakdown. The **ectonucleotidases** showed stereoselectivity towards the ribose moiety and towards the polyphosphate chain. In general, methylene isosteres of the nucleotides, and analogs in which 1 of the O atoms on the terminal phosphate had been replaced, were resistant to degradation. None of the analogs that were readily dephosphorylated was more potent than ATP, and most, but not all, of the analogs resistant to degradation were more potent than ATP, suggesting that although resistance to degradation does not in itself confer high potency, susceptibility to degradation does limit the potency of ATP and its degradable analogs.

IT 23567-96-6 23567-97-7 23600-16-0, 8-Bromo-ADP

RL: BIOL (Biological study)


(bladder contraction by, structure in relation to)

RN 23567-96-6 CAPLUS

CN 5'-Adenylic acid, 8-bromo- (9CI) (CA INDEX NAME)


*already posted
article of
these yds.*

Absolute stereochemistry.




RN 23567-97-7 CAPLUS
CN Adenosine 5'-(tetrahydrogen triphosphate), 8-bromo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 23600-16-0 CAPLUS
CN Adenosine 5'-(trihydrogen diphosphate), 8-bromo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 11 OF 11 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1987:12328 CAPLUS

DN 106:12328

TI ATP analogs and the guinea pig tenia coli: a comparison of the structure-activity relationships of **ectonucleotidases** with those of the P2-purinoceptor

AU Welford, Laurence A.; Cusack, Noel J.; Hourani, Susanna M. O.

CS King's Coll., Univ. London, London, WC2R 2LS, UK

SO European Journal of Pharmacology (1986), 129(3), 217-24

CODEN: EJPHAZ; ISSN: 0014-2999

DT Journal

LA English

AB The dephosphorylation of adenine nucleotides and their analogs by **ectonucleotidase** [9027-73-0] in the guinea pig tenia coli was studied using HPLC. The rate of dephosphorylation of each analog was compared with its pharmacol. potency relative to ATP [56-65-5]. ATP, ADP

*already printed
article 4 these copy.*

[58-64-0] and AMP [61-19-8] were rapidly dephosphorylated, and substitution on the purine ring had no effect upon the rate of breakdown. The **ectonucleotidases** showed stereoselectivity towards the ribose, the unnatural L-enantiomers of nucleotides being dephosphorylated more slowly. Analogs in which one of the O atoms on the terminal phosphate had been replaced were resistant to degradation. Phosphorothioate analogs in which a sulfur was attached to the penultimate phosphorus were degraded stereoselectively. Methylene isosteres of ATP and ADP resisted degradation, except for homo-ATP [72041-44-2] which was dephosphorylated at the same rate as ATP. Overall, no correlation was found between the potency of an analog and its rate of degradation.

IT 23567-96-6, 8-Bromoadenosine 5'-monophosphate 23567-97-7
 , 8-Bromoadenosine 5'-triphosphate 23600-16-0, 8-Bromoadenosine 5'-diphosphate


RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(metabolism of, by **ectonucleotidase** of *tenia coli*, P2-purinergic agonist activity in relation to)

RN 23567-96-6 CAPLUS

CN 5'-Adenylic acid, 8-bromo- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 23567-97-7 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-bromo- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 23600-16-0 CAPLUS

CN Adenosine 5'-(trihydrogen diphosphate), 8-bromo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



10/620,520

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal600txm

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

- NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
- NEWS 2 "Ask CAS" for self-help around the clock
- NEWS 3 JAN 17 Pre-1988 INPI data added to MARPAT
- NEWS 4 FEB 21 STN AnaVist, Version 1.1, lets you share your STN AnaVist visualization results
- NEWS 5 FEB 22 The IPC thesaurus added to additional patent databases on STN
- NEWS 6 FEB 22 Updates in EPFULL; IPC 8 enhancements added
- NEWS 7 FEB 27 New STN AnaVist pricing effective March 1, 2006
- NEWS 8 MAR 03 Updates in PATDPA; addition of IPC 8 data without attributes
- NEWS 9 MAR 22 EMBASE is now updated on a daily basis
- NEWS 10 APR 03 New IPC 8 fields and IPC thesaurus added to PATDPAFULL
- NEWS 11 APR 03 Bibliographic data updates resume; new IPC 8 fields and IPC thesaurus added in PCTFULL
- NEWS 12 APR 04 STN AnaVist \$500 visualization usage credit offered
- NEWS 13 APR 12 LINSPEC, learning database for INSPEC, reloaded and enhanced
- NEWS 14 APR 12 Improved structure highlighting in FQHIT and QHIT display in MARPAT
- NEWS 15 APR 12 Derwent World Patents Index to be reloaded and enhanced during second quarter; strategies may be affected
- NEWS 16 MAY 10 CA/CAPplus enhanced with 1900-1906 U.S. patent records
- NEWS 17 MAY 11 KOREAPAT updates resume
- NEWS 18 MAY 19 Derwent World Patents Index to be reloaded and enhanced
- NEWS EXPRESS FEBRUARY 15 CURRENT VERSION FOR WINDOWS IS V8.01a, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 19 DECEMBER 2005. V8.0 AND V8.01 USERS CAN OBTAIN THE UPGRADE TO V8.01a AT <http://download.cas.org/express/v8.0-Discover/>
- NEWS HOURS STN Operating Hours Plus Help Desk Availability
- NEWS LOGIN Welcome Banner and News Items
- NEWS IPC8 For general information regarding STN implementation of IPC 8
- NEWS X25 X.25 communication option no longer available after June 2006

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * *

COMPLETE THE STN SURVEY - APRIL 27 THROUGH MAY 31

Dear valued STN customer,

In an effort to enhance your experience with STN, we would like to better understand what you find useful. Please take approximately 5 minutes to complete a web survey.

If you provide us with your name, login ID, and e-mail address, you will be entered in a drawing to win a free iPod(R). Your responses will be kept confidential and will help us make future improvements to STN.

Take survey: <http://www.zoomerang.com/survey.zgi?p=WEB2259HNKWTUW>

Thank you in advance for your participation.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 00:34:05 ON 27 MAY 2006

=> d his

(FILE 'HOME' ENTERED AT 00:34:05 ON 27 MAY 2006)

=> file reg

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'REGISTRY' ENTERED AT 00:34:21 ON 27 MAY 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 25 MAY 2006 HIGHEST RN 885654-58-0

DICTIONARY FILE UPDATES: 25 MAY 2006 HIGHEST RN 885654-58-0

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 6, 2006

Please note that search-term pricing does apply when conducting SmartSELECT searches.

*
* The CA roles and document type information have been removed from *
* the IDE default display format and the ED field has been added, *
* effective March 20, 2005. A new display format, IDERL, is now *
* available and contains the CA role and document type information. *
*

Structure search iteration limits have been increased. See HELP SLIMITS for details.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10620520.str

L1 STRUCTURE UPLOADED

=> s l1 full

FULL SEARCH INITIATED 00:34:39 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 73256 TO ITERATE

100.0% PROCESSED 73256 ITERATIONS

1324 ANSWERS

SEARCH TIME: 00.00.01

L2 1324 SEA SSS FUL L1

=> file caplu

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

166.94

167.15

FILE 'CAPLUS' ENTERED AT 00:34:43 ON 27 MAY 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 27 May 2006 VOL 144 ISS 23

FILE LAST UPDATED: 25 May 2006 (20060525/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply.

They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> s aggregation

103702 AGGREGATION

2116 AGGREGATIONS

L3 105070 AGGREGATION

(AGGREGATION OR AGGREGATIONS)

=> s thrombogenicity

L4 2 THROMBOGENECITY

=> s thrombogenic?

L5 2 THROMBOGENEC?

=> s thrombogenic

=> s l2

L6 1225 L2

=> s l6 and (l3 or l4 or l5)

L7 14 L6 AND (L3 OR L4 OR L5)

=> d bib abs hitstr 1-14 17

L7 ANSWER 1 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2004:911398 CAPLUS

DN 142:214281

TI C8-substituted purine nucleotide analogs and their use as inhibitors of nucleoside triphosphate diphosphohydrolases

IN Halbfinger, Efrat; Fischer, Bilha; Beaudoin, Adrien R.; Gendron, Fernand Pierre

PA Universite de Sherbrooke, Can.; Bar-Ilan University

SO Can. Pat. Appl., 54 pp.

CODEN: CPXXEB

DT Patent

LA English

FAN.CNT 1

PATENT NO.

KIND

DATE

APPLICATION NO.


DATE

PI CA 2311084 AA 20011209 CA 2000-2311084 20000609
 PRAI CA 2000-2311084 20000609
 OS CASREACT 142:214281
 AB Ectonucleoside triphosphate diphosphohydrolases [NTPDases; EC 3.6.1.5] constitute a family of enzymes which are involved in the metabolism of extracellular nucleotides, catalyzing the hydrolysis of the gamma and beta phosphate bonds of triphospho- and diphosphonucleosides (whereas 5'-nucleotidases [EC 3.1.3.5] catalyze the hydrolysis of alpha phosphate bond of monophosphonucleosides). These extracellular nucleotides interact with endothelial, epithelial and smooth muscle cells, as well as blood cells and lymphoid cells, to influence the different physiol. systems of vertebrates. Since these ecto-nucleotidases alter the extracellular concns. of nucleotides these enzymes modulate their physiol. effects, including, for example, platelet **aggregation**, heart function, control of vascular tone and inflammation reactions, electrolyte secretion and gastrointestinal motility, neurotransmission both in central and peripheral nervous systems, as well as other effects in other physiol. systems. This invention provides C8 substituted purine nucleotide analogs, such as ATP analogs, and further provides their use as inhibitors of NTPDases and thereby as tools to modulate the conversion of nucleotides into nucleoside derivs., and thus modulate the levels of these compds. Such modulation further provides for the modulation of the activity and function of many processes which are affected by these compds.

IT 81609-35-0P 284040-51-3P 284040-52-4P
 284040-53-5P
 RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (C8-substituted purine nucleotide analogs and their use as inhibitors of nucleoside triphosphate diphosphohydrolases to modulate purine nucleotide levels and biol. processes)


RN 81609-35-0 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(ethylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 284040-51-3 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(cycloheptylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-52-4 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-[(2,2-dimethylpropyl)thio]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-53-5 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(hexylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 284040-54-6 284040-60-4

RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(C8-substituted purine nucleotide analogs and their use as inhibitors of nucleoside triphosphate diphosphohydrolases to modulate purine nucleotide levels and biol. processes)

RN 284040-54-6 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-60-4 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-butoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 23567-97-7, 8-Bromo-ATP

RL: BSU (Biological study, unclassified); BIOL (Biological study)
(substrate; C8-substituted purine nucleotide analogs and their use as
inhibitors of nucleoside triphosphate diphosphohydrolases to modulate
purine nucleotide levels and biol. processes)

RN 23567-97-7 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-bromo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L7 ANSWER 2 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1999:245290 CAPLUS


DN 131:59076

TI Structure-Activity Relationships of Bisphosphate Nucleotide Derivatives as
P2Y1 Receptor Antagonists and Partial Agonists

AU Nandan, Erathodiyil; Camaioni, Emidio; Jang, Soo-Yeon; Kim, Yong-Chul;
Cristalli, Gloria; Herdewijn, Piet; Secrist, John A., III; Tiwari, Kamal
N.; Mohanram, Arvind; Harden, T. Kendall; Boyer, Jose L.; Jacobson,
Kenneth A.

CS Molecular Recognition Section Laboratory of Bioorganic Chemistry, National
Institute of Diabetes Digestive and Kidney Diseases, National Institutes
of Health, Bethesda, MD, 20892-0810, USA

SO Journal of Medicinal Chemistry (1999), 42(9), 1625-1638



AB The P2Y1 receptor is present in the heart, in skeletal and various smooth muscles, and in platelets, where its activation is linked to **aggregation**. Adenosine 3',5'- and 2',5'-bis-phosphates have been identified as selective antagonists at the P2Y1 receptor and have been modified structurally to increase receptor affinity. We have extended the structure-activity relationships to a new series of deoxyadenosine bis-phosphates with substitutions in the adenine base, ribose moiety, and phosphate groups. The activity of each analog at P2Y1 receptors was determined by measuring its capacity to stimulate phospholipase C in turkey erythrocyte membranes (agonist effect) and to inhibit phospholipase C stimulation elicited by 10 nM 2-(methylthio)ADP (antagonist effect). 2'-Deoxyadenosine bis-phosphate analogs containing halo, amino, and thioether groups at the 2-position of the adenine ring were more potent P2Y1 receptor antagonists than analogs containing various heteroatom substitutions at the 8-position. An N6-methyl-2-chloro analog I, was a full antagonist and displayed an IC50 of 206 nM. On the ribose moiety, 2'-hydroxy, 4'-thio, carbocyclic, and six-membered anhydro-hexitol ring modifications have been prepared and resulted in enhanced agonist properties. The 1,5-anhydro-hexitol analog was a pure agonist with an EC50 of 3 μM, i.e., similar in potency to ATP 5'-Phosphate groups have been modified in the form of triphosphate, Me phosphate, and cyclic 3',5'-diphosphate derivs. The carbocyclic analog had enhanced agonist efficacy, and the 5'-O-phosphonyl-Me modification was tolerated, suggesting that deviations from the nucleotide structure may result in improved utility as pharmacol. probes. The N6-methoxy modification eliminated receptor affinity. Pyrimidine nucleoside 3',5'-bis-phosphate derivs. were inactive as agonists or antagonists at P2Y receptor subtypes.

IT 228264-34-4P 228264-35-5P 228264-36-6P
 228264-37-7P 228264-38-8P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(preparation and structure-activity relationships of bis-phosphate nucleotides as P2Y1 receptor antagonists and partial agonists)

RN 228264-34-4 CAPLUS

CN 3'-Adenylic acid, 2'-deoxy-8-methyl-, 5'-(dihydrogen phosphate), ammonium salt (9CI) (CA INDEX NAME)


Absolute stereochemistry.



●x NH₃

RN 228264-35-5 CAPLUS
 CN 3'-Adenylic acid, 2'-deoxy-8-ethenyl-, 5'-(dihydrogen phosphate), ammonium salt (9CI) (CA INDEX NAME)


Absolute stereochemistry.



●x NH₃

RN 228264-36-6 CAPLUS
 CN 3'-Adenylic acid, 2'-deoxy-8-methoxy-, 5'-(dihydrogen phosphate), ammonium salt (9CI) (CA INDEX NAME)


Absolute stereochemistry.



●x NH₃

RN 228264-37-7 CAPLUS
 CN 3'-Adenylic acid, 2'-deoxy-8-(methylthio)-, 5'-(dihydrogen phosphate), ammonium salt (9CI) (CA INDEX NAME)

Absolute stereochemistry.




●x NH₃

RN 228264-38-8 CAPLUS

CN 3'-Adenylic acid, 2'-deoxy-8-(methylamino)-, 5'-(dihydrogen phosphate), ammonium salt (9CI) (CA INDEX NAME)

Absolute stereochemistry.



●x NH₃

RE.CNT 39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 3 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1996:197574 CAPLUS

DN 124:313309

TI Platelet activation by 2-(4-bromo-2,3-dioxobutylthio)adenosine 5'-diphosphate is mediated by its binding to a putative ADP receptor, aggregin

AU Puri, Rajinder N.; Colman, Roberta F.; Colman, Robert W.

CS Sol Sherry Thrombosis Research Center, Temple Univ. School Medicine, Philadelphia, PA, USA

SO European Journal of Biochemistry (1996), 236(3), 862-70
CODEN: EJBCAI; ISSN: 0014-2956

PB Springer

DT Journal

LA English

AB Platelet responses induced by ADP are mediated by a unique P2T-purinergic receptor. Although a variety of ADP analogs, substituted at C2, were used to delineate pharmacol. properties of the ADP-binding site(s), the identity of the receptor protein was not firmly established. 2-(4-Bromo-2,3-dioxobutylthio)-ADP[2-BrCH₂(CO)₂CH₂-S-ADP], a well-characterized ADP analog, was used as an affinity label to examine the structure/function relationship of ADP-requiring enzymes. It induced platelet shape change, **aggregation**, exposure of fibrinogen binding sites, secretion, and mobilization of intracellular Ca, but was less potent than ADP. Under non-stirring conditions, incubation of platelets with this analog for longer time periods blocked ADP-induced shape change, **aggregation**, and the ability of ADP to antagonize

the rise in intracellular levels of cAMP induced by iloprost (a prostaglandin I2 analog). Of a variety of agonists examined, only ADP-induced **aggregation** was almost completely inhibited in platelets irreversibly modified by the analog. An autoradiogram of the gel obtained by SDS/PAGE of solubilized platelets modified by the ADP analog followed by reduction of the dioxo group by NaB[3H]4 showed the presence of a single radiolabeled protein band at 100 kDa. Platelets incubated 1st with either ADP, ATP, or 2-methylthio-ADP were not labeled by 2-BrCH2(CO)2CH2S-ADP and NaB[3H]4. 8-BrCH2(CO)2CH2S-ADP was previously shown by us to irreversibly antagonize ADP-induced platelet responses by selectively modifying aggregin. Incubation of platelets with 2-BrCH2(CO)2CH2S-ADP completely blocked labeling of aggregin in platelets by 8-BrCH2(CO)2CH2S-[32P]ADP. These results show that 2-BrCH2(CO)2CH2S-ADP initially interacts reversibly with aggregin (100 kDa), a putative ADP receptor, and induces platelet shape change and **aggregation**, and at longer periods of incubation reacts irreversibly to block the ability of ADP to antagonize stimulated adenylate cyclase activity. In contrast, 6-BrCH2(CO)2CH2S-ADP was found to be a weak and reversible inhibitor of ADP-induced platelet **aggregation**. Prior incubation of platelets with the latter analog reduced labeling of aggregin by 8-BrCH2(CO)2CH2S-[32P]ADP. Taken together, the results further show that substitution by the BrCH2(CO)2CH2 group at the C2 and C8 positions is tolerated, while the presence of a free amino function at the C6 position is essential for its interaction with aggregin.

IT 115678-78-9P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PNU (Preparation, unclassified); BIOL (Biological study); PREP (Preparation)

(platelet activation by 2-(4-bromo-2,3-dioxobutylthio)ADP is mediated by its binding to a putative ADP receptor, aggregin)

RN 115678-78-9 CAPLUS

CN Adenosine 5'-(trihydrogen diphosphate), 8-[(4-bromo-2,3-dioxobutyl)thio]-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L7 ANSWER 4 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1995:880408 CAPLUS

DN 123:311093

TI Inhibition of ADP-induced platelet responses by covalent modification of aggregin, a putative ADP receptor, by 8-(4-bromo-2,3-dioxobutylthio)ADP

AU Puri, Rajinder N.; Kumar, Ajay; Chen, Haiying; Colman, Roberta F.; Colman, Robert W.

CS Sol Sherry Thrombosis Res. Cent., Temple Univ. Sch. Med., Philadelphia, PA, 19140, USA

SO Journal of Biological Chemistry (1995), 270(41), 24482-8
CODEN: JBCHA3; ISSN: 0021-9258

PB American Society for Biochemistry and Molecular Bio logy

DT Journal

LA English

AB ADP is an important platelet agonist which initiates platelet shape change, **aggregation**, exposure of fibrinogen receptors, and calcium mobilization. Because of the limitations of previously used affinity analogs and photolabeling studies as well as controversies

surrounding the identity of an ADP receptor on platelets, we have used an affinity label capable of alkylating a putative exofacial receptor on platelets. We now report that 8-(4-bromo-2,3-dioxobutylthio)adenosine-5'-diphosphate (8-BDB-TADP), which is an analog of the natural ligand ADP, blocked ADP-induced platelet shape change, **aggregation**, exposure of fibrinogen-binding sites, secretion, and calcium mobilization. Following modification by 8-BDB-TADP, the rates of **aggregation** of platelets induced by thrombin, a calcium ionophore (A23187) or a stimulator of protein kinase C (phorbol myristate acetate) were minimally affected. However, the 8-BDB-TADP-modified platelets exhibited decreased rates of **aggregation** in response to ADP, as well as collagen and a thromboxane mimetic (U46619), both of which partially require ADP. Autoradiograms of the gels obtained by SDS-PAGE of solubilized platelets modified by either [β - 32 P]8-BDB-TADP, or 8-BDB-TADP and NaB[3 H]4 showed the presence of a single radiolabeled protein band at 100 kDa. The intensity of this band was reduced when platelets were preincubated with ADP, ATP, and 8-bromo-ADP prior to labeling by the radioactive 8-BDB-TADP. The results show that 8-BDB-TADP selectively and covalently labeled aggregin (100 kDa), a putative ADP receptor, resulting in a loss of ADP-induced platelet responses.

IT

115678-78-9

RL: ARG (Analytical reagent use); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)

(r; ADP-induced platelet responses inhibition by covalent modification of aggregin by (bromodioxobutylthio)ADP)


RN

115678-78-9 CAPLUS

CN

Adenosine 5'-(trihydrogen diphosphate), 8-[(4-bromo-2,3-dioxobutyl)thio]-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L7 ANSWER 5 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1994:695525 CAPLUS

DN 121:295525

TI Characterization of the HeLa Cell DNA Polymerase α -Associated Ap4A Binding Protein by Photoaffinity Labeling

AU Baxi, Mayur D.; McLennan, Alexander G.; Vishwanatha, Jamboor K.

CS Medical Center, University of Nebraska, Omaha, NE, 68198-4525, USA

SO Biochemistry (1994), 33(48), 14601-7

CODEN: BICHAW; ISSN: 0006-2960

PB American Chemical Society

DT Journal

LA English

AB The ubiquitous dinucleotide, diadenosine tetraphosphate (Ap4A), has been proposed to be involved in DNA replication and cell proliferation, DNA repair, platelet **aggregation**, and vascular tonus. A protein binding to Ap4A is associated with a multiprotein form of DNA polymerase- α (I) in HeLa cells. Here, the I-associated Ap4A-binding protein (II) was purified to homogeneity. II was resolved into 2 polypeptides of 45 and 22 kDa, designated A1 and A2, resp. [α - 32 P]8-azido (N3)-Ap4A was used to label purified II, and by crosslinking the photoaffinity label it was determined that Ap4A binds to the A1 subunit. No binding to the ligand was observed with the A2 subunit. Photoaffinity labeling was saturated with .apprx.0.4 μ M photolabel, with a

half-maximal binding at 0.15 μ M. The labeling was UV-dependent and was competed by both 8-N₃-Ap4A and Ap4A. Photoaffinity labeling was not affected by the presence of dATP and dGTP, and was reduced only in the presence of excess of ATP, indicating the specificity of II for Ap4A. Of the diadenosine polyphosphates, Ap4A and Ap5A competed for binding, whereas Ap2A and Ap3A did not compete for binding. Further, the presence of ≥ 1 adenosine(s) may be necessary since Ap4G competed but Gp4G did not compete for binding to II. Various methylene bisphosphonate and thiophosphate analogs of Ap4A were tested for their effect on photoaffinity labeling with 8-N₃-Ap4A. Significant differences were observed among the various analogs in their ability to prevent the photoaffinity labeling of the ligand to II.

IT 126813-90-9, 8-Azido-Ap4A


RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(purification and photoaffinity labeling of HeLa cell DNA polymerase- α -associated Ap4A-binding protein)

RN 126813-90-9 CAPLUS


CN Adenosine 5'-(pentahydrogen tetraphosphate), 8-azido-,
P''' \rightarrow 5'-ester with adenosine (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B



L7 ANSWER 6 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1994:401198 CAPLUS

DN 121:1198

TI Interaction of Nucleotides with Acidic Fibroblast Growth Factor (FGF-1)
AU Chavan, Ashok J.; Haley, Boyd E.; Volkin, David B.; Marfia, Kimberly E.; Verticelli, Adeline M.; Bruner, Mark W.; Draper, Jerome P.; Burke, Carl J.; Middaugh, C. Russell

CS College of Pharmacy, University of Kentucky, Lexington, KY, 40536, USA

SO Biochemistry (1994), 33(23), 7193-202

CODEN: BICHAW; ISSN: 0006-2960

DT Journal

LA English

AB A wide variety of nucleotides is shown to bind to acidic fibroblast growth factor (aFGF) as demonstrated by their ability to inhibit the heat-induced aggregation of the protein; to enhance the thermal stability of aFGF as monitored by both intrinsic fluorescence and CD; to interact with fluorescent nucleotides and displace a bound polysulfated naphthylurea compound, suramin; to reduce the size of heparin-aFGF complexes; and to protect a reactive aFGF thiol group. The binding of mononucleotides, diadenosine compds. (ApnA), and inorg. polyphosphates to aFGF is enhanced as the degree of phosphorylation of these anions is increased with the presence of the base reducing the apparent binding affinity. The nature of the base appears to have much less effect. Photoactivatable nucleotides (8N3-ATP, 2N3-ATP, 8N3-GTP, and 8N3-Ap4A) were employed to covalently label the aFGF nucleotide binding site. In general, Kd's in the low micromolar range are observed. Protection against 90% displacement is observed at several hundred micromolar nucleotide concentration. Using 8N3-ATP as a prototypic reagent, photolabeled aFGF was proteolyzed with trypsin and chymotrypsin, and labeled peptides were isolated and sequenced resulting in the identification of 10 possible labeled amino acids (Y8, G20, H21, T61, K112, K113, S116, R119, R122, H124). On the basis of the crystal structure of bovine aFGF, eight of the prospective labeled sites appear to be dispersed around the perimeter of the growth factor's presumptive polyanion binding site. One residue (T61) is more distally located but still proximate to several pos. charged residues, and another (Y8) is not locatable in crystal structures. Using heparin affinity chromatog., at least three distinct photolabeled aFGF species were resolved. These labeled complexes display diminished affinity for heparin and a reduced ability to stimulate mitogenesis even in the presence of polyanions such as heparin. In conclusion, nucleotides bind apparently nonspecifically to the polyanion binding site of aFGF but nevertheless are capable of modulating the protein's activity. Evidence for the presence of a second or more extended polyanion binding site and the potential biol. significance of these results in terms of potential natural ligands of aFGF are also discussed but not resolved.

IT 53696-59-6, 8-Azido-ATP 65114-35-4, 8-Azido-GTP

126813-90-9, 8-Azido-Ap4A


RL: BIOL (Biological study)

(acidic fibroblast growth factor binding of, sites for)

RN 53696-59-6 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-azido- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 65114-35-4 CAPLUS

CN Guanosine 5'-(tetrahydrogen triphosphate), 8-azido- (9CI) (CA INDEX NAME)


Absolute stereochemistry.




RN 126813-90-9 CAPLUS
 CN Adenosine 5'-(pentahydrogen tetraphosphate), 8-azido-,
 P'''->5'-ester with adenosine (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B



L7 ANSWER 7 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 1993:490701 CAPLUS
 DN 119:90701
 TI Identification of a receptor for ADP on blood platelets by photoaffinity labeling
 AU Cristalli, Gloria; Mills, David C. B.
 CS Sch. Med., Temple Univ., Philadelphia, PA, 19140, USA
 SO Biochemical Journal (1993), 291(3), 875-81
 CODEN: BIJOAK; ISSN: 0306-3275
 DT Journal
 LA English
 AB The synthesis of a new analog of ADP, 2-(p-azidophenyl)-ethylthioadenosine 5'-diphosphate (AzPET-ADP), is described. This compound contains a photolabile phenylazide group attached to the ADP mol. by a thioether link at the purine 2 position. It has been prepared in radioactive form with ³²P

in the β -phosphate at a specific radioactivity of 100 mCi/ μ mol. The reagent activated platelets, causing shape change and **aggregation**, with somewhat lower affinity than ADP. On photolysis the affinity was increased. The reagent also inhibited platelet adenylate cyclase stimulation by prostaglandin E₁, with considerably higher affinity than ADP. On photolysis the affinity was decreased. AzPET-ADP competitively inhibited the binding of 2-methylthio[β -³²P]ADP, a ligand for the receptor by which ADP causes inhibition of adenylate cyclase. In the dark, AzPET-[β -³²P]ADP bound reversibly and with high affinity to a single population of sites similar in number to the sites that bind 2-methylthiol[β -³²P]ADP. Binding was inhibited by ADP and by ATP and by p-chloromercuribenzenesulfonic acid (pCMBS). On exposure to UV light in the presence of platelets, AzPET-[β -³²P]ADP was incorporated covalently but non-specifically into several platelet proteins, although prominent intracellular proteins were not labeled. Specific labeling was confined to a single region of SDS/polyacrylamide gels, overlying but not comigrating with actin. Incorporation of radioactivity into this region was inhibited by ADP and by ATP as well as by ADP β S, ATP α S and pCMBS, but not by adenosine, GDP or AMP. Inhibition of AzPET-[β -³²P]ADP incorporation was closely correlated with inhibition of equilibrium binding of 2-methylthio[β -³²P]ADP. These results suggest that the labeled protein, which migrates with an apparent mol. mass of 43 kDa in reduced gels, is the receptor through which ADP inhibits adenylate cyclase.

IT 23600-16-0


RL: ANST (Analytical study)

(binding of azidophenylethylthioadenosine diphosphate and methylthio ADP into protein in blood platelets in relation to)

RN 23600-16-0 CAPLUS

CN Adenosine 5'-(trihydrogen diphosphate), 8-bromo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L7 ANSWER 8 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1991:464361 CAPLUS

DN 115:64361

TI The effect of agonists and antagonists of platelet **aggregation** on von Willebrand factor-mediated platelet agglutination

AU McPherson, Jean; Zucker, Marjorie B.; Mauss, Evelyn A.; Brownlea, Sandra
CS Fac. Med., Univ. Newcastle, Newcastle, 2308, Australia

SO Thrombosis and Haemostasis (1991), 65(5), 573-7

CODEN: THHADQ; ISSN: 0340-6245

DT Journal

LA English

AB Ristocetin-induced platelet agglutination (RIPA) in EDTA-treated citrated platelet-rich plasma was reduced to 49% by 1.25 μ M ADP, 41% by 1 μ M A23187, and 26% by 0.1 μ g/mL platelet activating factor (PAF). The effect of 5-110 μ M epinephrine was not dose-dependent, but varied between donors, with RIPA from 56-100% of the control. The inhibitory effects of these agonists were not altered by prior treatment of platelets with aspirin. Prior addition of 200 μ M ATP (an ADP receptor antagonist acting at both high- and low-affinity ADP receptors) prevented the inhibitory action of ADP but not that of A23187 or PAF, suggesting that the inhibitory actions of the latter were not mediated by released ADP. As 700 μ M 8-bromoadenosine 5-diphosphate (an ADP receptor antagonist


acting mainly at the high-affinity receptor) did not prevent the ADP-induced inhibition of RIPA, the interaction of ADP with the low-affinity receptor was presumably responsible for its inhibitory action. As A23187, but not phorbol myristate acetate (0.1 μ M), inhibited RIPA, an increase in intracellular calcium ions rather than direct stimulation of protein kinase C appears to mediate the agonist-induced inhibition. Cytochalasin B (10.5-21 μ M), dibucaine (0.5-1 mM), and PGE1 (25 nM) added before or after the agonist prevented or reversed the ADP-, A23187-, and PAF-induced inhibition of RIPA, suggesting that the state of the platelet cytoskeleton affects the inhibition. N-Ethylmaleimide (0.25-0.5 mM), an agent that can penetrate cell membranes and block sulfhydryl groups, prevented or reversed the ADP-, A23187- and PAF-induced inhibition of RIPA, but 0.5 mM dithionitrobenzoic acid, a non-penetrating sulfhydryl blocker, had no effect. Diamide (0.1-0.5 mM), an agent that can crosslink cytoskeletal proteins by oxidation of sulfhydryl groups, reduced RIPA. Thus, an increase in intracellular calcium ions with resultant cytoskeletal changes and reorganization of intracellular sulfhydryl groups may mediate the inhibitory action of agonists on RIPA.

IT 23600-16-0, 8-Bromo-ADP
 RL: BIOL (Biological study)
 (blood platelet **aggregation** induced by ristocetin response to)

RN 23600-16-0 CAPLUS

CN Adenosine 5'-(trihydrogen diphosphate), 8-bromo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L7 ANSWER 9 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1989:511319 CAPLUS

DN 111:111319

TI The role of nucleotide self-**aggregation** on the interaction process between clupeine YI (a fish protamine) and mono- and dinucleotides

AU Andini, Salvatore; Ferrara, Luciano; Cocozziello, Beatrice; De Napoli, Lorenzo; Piccialli, Gennaro; Barbato, Stefania

CS Dip. Chim., Univ. Napoli, Naples, I-80134, Italy

SO Gazzetta Chimica Italiana (1989), 119(5), 271-5

CODEN: GCITA9; ISSN: 0016-5603

DT Journal

LA English

AB The interaction between clupeine YI and various nucleotides in aqueous solution has been investigated with the aim of understanding the reason for the stronger affinity of purinic nucleotides towards the protein as compared with pyrimidinic ones. The study has been performed by ¹H NMR spectrometry by adding increasing amts. of nucleotide to the protamine solution. The results obtained suggest that it is not the nature (purinic or pyrimidinic) of the nucleotide but its ability to give rise to a self-**aggregation** process that is crucial in the interaction with the protein. In fact, nucleotide complexation should provide a polyanionic matrix around which the protein can be strongly linked. This model was tested with different natural, synthetic, and modified nucleotides and with some dinucleotides.

IT 21870-09-7, 8-Bromo-5'-gmp 23567-96-6, 8-Bromo-5'-amp
 61286-93-9


RL: BIOL (Biological study)

(protamine clupeine Y1 interaction with, nucleotide self-
aggregation role in)

RN 21870-09-7 CAPLUS

CN 5'-Guanylic acid, 8-bromo- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 23567-96-6 CAPLUS

CN 5'-Adenylic acid, 8-bromo- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 61286-93-9 CAPLUS

CN 5'-Adenylic acid, 8-bromo-2'-deoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L7 ANSWER 10 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1986:437246 CAPLUS

DN 105:37246

TI Proton NMR study of self-association and restricted internal rotation of
the C8-substituted deoxyguanosine 5'-monophosphate adduct of the
carcinogen 2-(acetylamino)fluorene

AU Evans, Frederick E.; Miller, Dwight W.; Levine, Robert A.


CS Div. Chem., Natl. Cent. Toxicol. Res., Jefferson, AR, 72079, USA

SO Journal of Biomolecular Structure & Dynamics (1986), 3(5), 935-48
CODEN: JBSDD6; ISSN: 0739-1102

DT Journal

LA English

GI



AB The high-field ^1H NMR spectra of a nucleotide-carcinogen adduct formed from 2-(acetylamino)fluorene, [8-(N-fluorene-2-ylacetamido)-2'-deoxyguanosine 5'-monophosphate (I) [14490-86-9]], was examined in aqueous solution as a function of concentration at high and low temps. An anomalous concentration dependence of NMR spectra was observed at >1 mM. These spectral characteristics were analyzed in terms of changes in selfassocn. and in the interconversions between torsional diastereomers associated with the central N. Association consts. were computed. Stacking interactions, which involve the fluorene and guanine rings, are strong, cooperative and highly temperature-dependent. Deacetylation alters the mode of stacking. Several effects of solvent and aggregation on the conformation at the central N are discussed.


IT 14490-86-9

RL: BIOL (Biological study)
(NMR study of structure of)

RN 14490-86-9 CAPLUS

CN 5'-Guanylic acid, 8-(acetyl-9H-fluorene-2-ylamino)-2'-deoxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L7 ANSWER 11 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1986:84344 CAPLUS

DN 104:84344

TI Distances between active site probes in glutamine synthetase from Escherichia coli: fluorescence energy transfer in free and in stacked dodecamers

AU Maurizi, Michael R.; Kasprzyk, Philip G.; Ginsburg, Ann

CS Lab. Mol. Biol., Natl. Cancer Inst., Bethesda, MD, 20205, USA

SO Biochemistry (1986), 25(1), 141-51

CODEN: BICHAW; ISSN: 0006-2960

DT Journal

LA English


AB Probes for fluorescence energy transfer measurement were introduced into active sites of dodecameric glutamine synthetase from *Escherichia coli* by substituting appropriate ATP analogs for ATP in the autoinactivation reaction of this enzyme with L-methionine-(S)-sulfoximide and Mn^{2+} . Two fluorescent donors, 8-mercapto-ATP alkylated with either 5-[[[(iodoacetyl)amino]ethyl]amino]naphthalene-1-sulfonic acid (AEDANS-ATP) or 1,N6-etheno-2-aza-ATP (aza- ϵ -ATP), and 2 acceptors, 6-mercaptapurine ribonucleotide triphosphate or 8-mercapto-ATP alkylated with the chromophore [[(4-dimethylamino)phenyl]azo]-2-iodoanilide (6-Y- or 8-Y-ATP), were used. Fluorescence emissions of enzyme derivs. with 1 or 2 equivalent of fluorescent donor/dodecamer and either an acceptor (Y) or ADP at the remaining active sites were compared at pH 7.0. The results, together with the known geometry of the enzyme, indicate that active-site probes in the dodecamer are widely separated and that energy transfer occurs from a single donor to 2 or 3 acceptors on adjacent subunits. The calculated distance between equidistant active-site probes on heterologously bonded subunit within the same hexagonal ring is 56-60 Å. Probes on isologously bonded subunits are no closer than 60 Å and may be as far apart as 78 Å. Thus, active sites are away from the 6-fold axis of symmetry toward the outer edges of the dodecamer and are located ≥ 30 Å from the plane separating the hexagonal rings. During Zn^{2+} -induced stacking of the same enzyme derivs. along the 6-fold axes of symmetry, addnl. quenches of fluorescence probes were dependent on the presence of acceptors on sep. dodecamers. The Zn^{2+} -induced face-to-face **aggregation** of dodecamers in the presence of 46 μM $ZnCl_2$ and 9 mM $MgCl_2$ at pH 7.0 and an Arrhenius activation energy of 22.3 kcal/mol and a 2nd-order rate constant at 25° of .apprx.105 M⁻¹ s⁻¹ at early stages. Time-dependent fluorescence quenches maximum values of 47-70% quench when the average oligomer was dodecamers. After correction for unquenched polymer ends, a fluorescent donor and an acceptor probe in layered dodecamers were estimated to be .apprx.36 Å apart (an average value if there is some twisting of single strands). This intermol. energy-transfer distance confirms that activity-site nucleotide probes are located toward exterior surfaces away from the lateral plane separating hexagonal rings of a dodecamer.

IT **99376-89-3**
 RL: BIOL (Biological study)
 (fluorescence energy transfer to acceptor from, in glutamine synthetase of *Escherichia coli*)

RN 99376-89-3 CAPLUS

CN 1-Naphthalenesulfonic acid, 5-[[[2-[[[[[6-amino-9-[5-O-[hydroxy[[hydroxy(phosphonooxy)phosphinyl]oxy]phosphinyl]- β -D-ribofuranosyl]-9H-purin-8-yl]thio]acetyl]amino]ethyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT **99376-89-3D**, reaction products with glutamine synthetase


RL: PRP (Properties)

(fluorescence of)

RN 99376-89-3 CAPLUS

CN 1-Naphthalenesulfonic acid, 5-[[2-[[[6-amino-9-[5-O-[hydroxy[[hydroxy(phosphonooxy)phosphinyl]oxy]phosphinyl]-β-D-ribofuranosyl]-9H-purin-8-yl]thio]acetyl]amino]ethyl]amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT **99376-91-7P**

RL: SPN (Synthetic preparation); PREP (Preparation)


(preparation and use in glutamine synthetase active site of Escherichia coli anal.)

RN 99376-91-7 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-[[2-[[4-[[4-


(dimethylamino)phenyl]azo]phenyl]amino]-2-oxoethyl]thio]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry unknown.



IT 41106-66-5
RL: RCT (Reactant); RACT (Reactant or reagent)
(reaction of, with iodoacetanilide derivative)
RN 41106-66-5 CAPLUS
CN Adenosine 5'-(tetrahydrogen triphosphate), 7,8-dihydro-8-thioxo- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.



L7 ANSWER 12 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN
AN 1983:519886 CAPLUS
DN 99:119886
TI Interactions of two ADP analogs, xanthosine-5'-diphosphate and 8-bromoadenosine-5'-diphosphate, and ADP with citrated human platelet-rich plasma
AU Ragatz, Barth H.
CS Sch. Med., Indiana Univ., Fort Wayne, IN, 46805, USA
SO Proceedings of the Indiana Academy of Science (1982), Volume Date 1981, 91, 183-7
CODEN: PIACAP; ISSN: 0073-6767
DT Journal
LA English
AB To determine the stereochem. nature of the ADP receptor of human platelets, the interaction of 2 ADP analogs (xanthosine 5'-diphosphate (XDP) and 8-bromo-ADP) with human platelet-rich plasma was investigated. XDP over a 100-fold concentration range failed to yield any competitive inhibition of ADP-induced platelet **aggregation**. Completely analogous results were observed for bromo-ADP. When either compound was preincubated with the platelet-rich plasma for 5 min prior to ADP addition, no inhibition of

ADP-induced **aggregation** occurred. Thus, substituents at position C-2 and C-6 (as seen with XDP) and C-8 (bromo-ADP) are not tolerated at the ADP receptor.

IT 23600-16-0


RL: BIOL (Biological study)

(ADP receptors stereochem. in human blood platelet in relation to)

RN 23600-16-0 CAPLUS

CN Adenosine 5'-(trihydrogen diphosphate), 8-bromo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L7 ANSWER 13 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1980:582136 CAPLUS

DN 93:182136

TI Affinity chromatography on immobilized nucleotides. The synthesis, specificity and applications of immobilized inosine 5'-monophosphate

AU Clonis, Yannis D.; Lowe, Christopher R.

CS Dep. Biochem., Univ. Southampton, Southampton, UK

SO European Journal of Biochemistry (1980), 110(1), 279-88

CODEN: EJBCAI; ISSN: 0014-2956

DT Journal

LA English

AB The synthesis and characterization of 2 IMP analogs, 8-(6-aminohexyl)-IMP and inosine 2',3'-O-[1-(6-aminohexyl)-levulinic acid amide]-acetal 5'-monophosphate are described. These analogs were attached to CNBr-activated agarose through the terminal NH₂ group of the spacer mol. The immobilized IMP analogs displayed specificity for the inosine-nucleotide-dependent enzyme IMP dehydrogenase but not for the NAD⁺-dependent enzymes L-alanine and L-acetate dehydrogenases. Escherichia coli IMP dehydrogenase could be eluted biospecifically from immobilized 8-substituted and ribose-substituted IMP adsorbents with IMP, XMP, and GMP. Multiple peaks of enzyme activity in the elution profiles were interpreted in terms of **aggregation** of the enzyme. A protocol for the large-scale purification of E. coli IMP dehydrogenase is proposed. Homogeneous enzyme of sp. activity 9.1 units/mg was obtained in 50% overall yield, representing 14 mg pure protein from a 20-L culture of E. coli. The 2 IMP analogs were inactive as substrates in the IMP dehydrogenase reaction.

IT 75204-34-1P


RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation and immobilization of, on Sepharose for affinity chromatog.)

RN 75204-34-1 CAPLUS

CN 5'-Inosinic acid, 8-[(6-aminohexyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 75204-33-0P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and reaction of, with diamino-hexane)

RN 75204-33-0 CAPLUS

CN 5'-Inosinic acid, 8-bromo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 52977-32-9DP, reaction products with Sepharose 4B

RL: PREP (Preparation)
(preparation of, for affinity chromatog.)

RN 52977-32-9 CAPLUS

CN 5'-Adenylic acid, 8-[(6-amino-hexyl)amino]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L7 ANSWER 14 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1976:472930 CAPLUS

DN 85:72930

TI Sulfhydryl analogs of adenosine diphosphate: chemical synthesis and activity as platelet-aggregating agents

AU Stone, J. V.; Singh, Raj K.; Horak, H.; Barton, P. G.

CS Dep. Biochem., Univ. Alberta, Edmonton, AB, Can.


SO Canadian Journal of Biochemistry (1976), 54(6), 529-33

CODEN: CJBIAE; ISSN: 0008-4018

DT Journal

LA English

GI



AB 2-Thioadenosine 5'-diphosphate (2-SH ADP) (I) [59924-53-7],
 2,2'-dithiobisadenosine 5'-diphosphate (2,2'-(S-ADP)2) [59924-54-8],
 8-thioadenosine 5'-diphosphate triethylammonium salt (8SH ADP) [59924-56-0], and 6-mercaptapurineriboside 5'-disphosphate (6-MPRDP) [805-63-0] were synthesized as potential affinity labels for ADP receptors on the blood-platelet membrane. The mean relative activities of these compds. in aggregating human platelets suspended in homologous plasma were 155% (2,2'-(S-ADP)2), 74% (2-SH ADP), 0.65% (8-SH ADP), and 0.08% (6-MPRDP). The mean relative activities against washed platelets were 249% (2,2'-(S-ADP)2) and 115% (2-SHADP), whereas no **aggregation** occurred with 8-SH ADP or 6-MPRDP. The last 2 compds. were weak inhibitors of ADP-induced **aggregation**. Therefore, thio-substitution at position 2 followed by oxidation to a disulfide appears to be the most promising approach to further studies of affinity labeling of membrane ADP-receptors.

IT 59924-56-0P

RL: PREP (Preparation)

(preparation and blood platelet **aggregation** response to)

RN 59924-56-0 CAPLUS


CN Adenosine 5'-(trihydrogen diphosphate), 7,8-dihydro-8-thioxo-, compd. with N,N-diethylethanamine (1:1) (9CI) (CA INDEX NAME)

CM 1

CRN 59924-55-9

CMF C10 H15 N5 O10 P2 S


Absolute stereochemistry.



CM 2


CRN 121-44-8

CMF C6 H15 N



IT 34051-09-7P
 RL: PREP (Preparation)
 (preparation of)
 RN 34051-09-7 CAPLUS
 CN 5'-Adenylic acid, 7,8-dihydro-8-thioxo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



=> FIL STNGUIDE		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	79.69	246.84
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-10.50	-10.50

FILE 'STNGUIDE' ENTERED AT 00:37:25 ON 27 MAY 2006
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
 COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE
 AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.
 LAST RELOADED: May 19, 2006 (20060519/UP).

10/620,520

(FILE 'HOME' ENTERED AT 08:47:16 ON 13 MAY 2006)

FILE 'REGISTRY' ENTERED AT 08:47:22 ON 13 MAY 2006

L1 STRUCTURE UPLOADED
L2 20 S L1 SSS SAM
L3 STRUCTURE UPLOADED
L4 19 S L3 SSS SAM

FILE 'REGISTRY' ENTERED AT 09:08:57 ON 13 MAY 2006

L5 STRUCTURE UPLOADED
L6 2 S L5 SSS SAM
L7 32 S L5 SSS FULL

FILE 'CAPLUS' ENTERED AT 09:10:18 ON 13 MAY 2006

L8 17 S L7

=> s l8 and ntpdase
112 NTPDASE
47 NTPDASES
128 NTPDASE
(NTPDASE OR NTPDASES)
L9 5 L8 AND NTPDASE

=> d bib abs hitstr 1-5 l9

L9 ANSWER 1 OF 5 CAPLUS COPYRIGHT 2006 ACS on STN
AN 2004:911398 CAPLUS
DN 142:214281
TI C8-substituted purine nucleotide analogs and their use as inhibitors of
nucleoside triphosphate diphosphohydrolases
IN Halbfinger, Efrat; Fischer, Bilha; Beaudoin, Adrien R.; Gendron, Fernand
Pierre
PA Universite de Sherbrooke, Can.; Bar-Ilan University
SO Can. Pat. Appl., 54 pp.
CODEN: CPXXEB
DT Patent
LA English
FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	---	-----	-----	-----
PI	CA 2311084	AA	20011209	CA 2000-2311084	20000609
PRAI	CA 2000-2311084		20000609		
OS	CASREACT 142:214281				
AB	Ectonucleoside triphosphate diphosphohydrolases [NTPDases; EC 3.6.1.5] constitute a family of enzymes which are involved in the metabolism of extracellular nucleotides, catalyzing the hydrolysis of the gamma and beta phosphate bonds of triphospho- and diphosphonucleosides (whereas 5'-nucleotidases [EC 3.1.3.5] catalyze the hydrolysis of alpha phosphate bond of monophosphonucleosides). These extracellular nucleotides interact with endothelial, epithelial and smooth muscle cells, as well as blood cells and lymphoid cells, to influence the different physiol. systems of vertebrates. Since these ecto-nucleotidases alter the extracellular concns. of nucleotides these enzymes modulate their physiol. effects, including, for example, platelet aggregation, heart function, control of vascular tone and inflammation reactions, electrolyte secretion and gastrointestinal motility, neurotransmission both in central and peripheral nervous systems, as well as other effects in other physiol. systems. This invention provides C8 substituted purine nucleotide analogs, such as ATP analogs, and further provides their use as inhibitors of NTPDases and thereby as tools to modulate the conversion of nucleotides into nucleoside derivs., and thus modulate the levels of these compds. Such modulation further provides for the modulation of the activity and function of many processes which are affected by these compds.				
IT	81609-35-0P 284040-51-3P 284040-52-4P 284040-53-5P RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)				

(C8-substituted purine nucleotide analogs and their use as inhibitors of nucleoside triphosphate diphosphohydrolases to modulate purine nucleotide levels and biol. processes)

RN 81609-35-0 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(ethylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-51-3 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(cycloheptylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-52-4 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-[(2,2-dimethylpropyl)thio]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-53-5 CAPLUS


CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(hexylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 284040-54-6
 RL: BSU (Biological study, unclassified); PAC (Pharmacological activity);
 THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (C8-substituted purine nucleotide analogs and their use as inhibitors
 of nucleoside triphosphate diphosphohydrolases to modulate purine
 nucleotide levels and biol. processes)
 RN 284040-54-6 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX
 NAME)

Absolute stereochemistry.



L9 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2003:856092 CAPLUS
 DN 139:333119
 TI Ecto-nucleoside triphosphate diphosphohydrolase inhibition-based methods
 for screening for a compound useful in the treatment or prevention of
 lymphocytic disorders, for inhibiting lymphocyte activity and preventing
 or treating lymphocytic disorders
 IN Beaudoin, Adrien; Benrezzak, Ouhida
 PA Bioflash Inc., Can.
 SO PCT Int. Appl., 63 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003089664	A1	20031030	WO 2003-CA583	20030422
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	CA 2382768	AA	20031019	CA 2002-2382768	20020419
	CA 2479501	AA	20031030	CA 2003-2479501	20030422


AU 2003226989	A1	20031103	AU 2003-226989	20030422
US 2005164306	A1	20050728	US 2003-511133	20030422
PRAI CA 2002-2382768	A	20020419		
WO 2003-CA583	W	20030422		

AB The invention discloses a method of screening for a compound useful in the treatment of a disease or condition characterized by an immune cell disorder, wherein the cell expresses ecto-nucleoside triphosphate diphosphohydrolases (**NTPDases**), the method comprising contacting a candidate compound with **NTPDase**, wherein the candidate compound is selected if the activity of the **NTPDase** is reduced in the presence of the candidate compound as compared to that in the absence thereof. The invention also discloses a method for inhibiting an immune cell activity in a mammal, comprising targeting immune cells with an effective amount of a **NTPDase** inhibitor. The invention further discloses a method to prevent or reduce the risk of rejection of transplanted tissue or organ, comprising administering to the animal an effective amount of **NTPDase** inhibitor.

IT 284040-54-6 344402-39-7
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (ecto-nucleoside triphosphate diphosphohydrolase inhibition-based methods for screening for agents for treatment of immune cell disorder-associated conditions)


RN 284040-54-6 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 344402-39-7 CAPLUS
 CN 5'-Adenylic acid, 8-(butylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 3 OF 5 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2003:711173 CAPLUS
 DN 139:230955
 TI Preparation of C8-substituted purine nucleotide analogs as **NTPDase** inhibitors
 IN Beaudoin, Adrien R.; Gendron, Fernand-Pierre; Halbfinger, Efrat; Fischer,

PA Bilha
 SO Bar-Ilan University, Israel; Universite De Sherbrooke
 U.S., 21 pp.
 CODEN: USXXAM
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 6617439	B1	20030909	US 2000-591177	20000609
	US 2004043955	A1	20040304	US 2003-620520	20030716
PRAI	US 2000-591177	A3	20000609		
OS	MARPAT 139:230955				
GI					

Parent



AB C8-substituted purine nucleotide analogs, I (R is alkyl, cycloalkyl) such as ATP analogs, and their use is described, including their use as inhibitors of **NTPDases** and thereby as tools to modulate the conversion of nucleotides into nucleoside derivs., and thus modulate the levels of these compds. Such modulation further provides for the modulation of the activity and function of many processes which are affected by these compds. Thus, I [R = (CH₂)₃Me] was prepared and tested in vivo as NTPDase inhibitor. I [R = (CH₂)₃Me] interacts specifically with the binding site of the enzyme potentially reduces the risk of interference with other ATP-binding enzymes or receptors, and thus possesses a high degree of specificity. The compds. of the invention were analyzed with resp. to any effects on the activity of purinoceptors.

IT 81609-35-0P 284040-51-3P 284040-52-4P
 284040-53-5P 284040-54-6P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of C8-substituted purine nucleotide analogs as **NTPDase** inhibitors)

RN 81609-35-0 CAPLUS


CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(ethylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 284040-51-3 CAPLUS
CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(cycloheptylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 284040-52-4 CAPLUS
CN Adenosine 5'-(tetrahydrogen triphosphate), 8-[(2,2-dimethylpropyl)thio]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 284040-53-5 CAPLUS
CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(hexylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 284040-54-6 CAPLUS
CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 71 THERE ARE 71 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 4 OF 5 CAPLUS COPYRIGHT 2006 ACS on STN
AN 2001:327844 CAPLUS
DN 135:149038
TI Inhibitors of **NTPDase**: key players in the metabolism of
extracellular purines
AU Gendron, F. P.; Halbfinger, E.; Fischer, B.; Beaudoin, A. R.
CS Department of Biology, University of Sherbrooke, Sherbrooke, Can.
SO Advances in Experimental Medicine and Biology (2000), 486(Purine and
Pyrimidine Metabolism in Man X), 119-123
CODEN: AEMBAP; ISSN: 0065-2598

PB Kluwer Academic/Plenum Publishers
DT Journal
LA English

AB This study described the potential of a new class of ATP analogs as
nucleoside triphosphate diphosphohydrolase (**NTPDase**) inhibitors.
From previous studies, 8-thiobutyladenosine 5'-triphosphate (8-BuS-ATP)
appears to be a specific and efficient **NTPDase** inhibitor. This
novel inhibitor is a new tool to regulate **NTPDase** activity and
thereby influencing purine signaling in mammalian.


IT 81609-35-0 284040-51-3 284040-53-5
284040-54-6

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL
(Biological study); PROC (Process)
(inhibitors of nucleoside triphosphate diphosphohydrolase - key players
in metabolism of extracellular purines)

RN 81609-35-0 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(ethylthio)- (9CI) (CA INDEX
NAME)

Absolute stereochemistry.




RN 284040-51-3 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(cycloheptylthio)- (9CI) (CA
INDEX NAME)


Absolute stereochemistry.

Adonis
Printed




RN 284040-53-5 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(hexylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 284040-54-6 CAPLUS
 CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 5 OF 5 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2000:304989 CAPLUS
 DN 133:105244
 TI Novel Inhibitors of Nucleoside Triphosphate Diphosphohydrolases: Chemical Synthesis and Biochemical and Pharmacological Characterizations
 AU Gendron, Fernand-Pierre; Halbfinger, Efrat; Fischer, Bilha; Duval, Martine; D'Orleans-Juste, Pedro; Beaudoin, Adrien R.
 CS Department de Biologie, Universite de Sherbrooke, Sherbrooke, QC, J1K 2R1, Can.
 SO Journal of Medicinal Chemistry (2000), 43(11), 2239-2247
 CODEN: JMCMAR; ISSN: 0022-2623

Printed

PB American Chemical Society

DT Journal

LA English

AB To elucidate the physiologic role played by nucleoside triphosphate diphosphohydrolase (**NTPDase**; EC 3.6.1.5), adenine nucleotide analogs, modified on the purine ring, have been synthesized and tested as potential inhibitors. Resistance of ATP analogs to hydrolysis and their potency as **NTPDase** inhibitors were evaluated. For this purpose, a particulate fraction isolated from bovine spleen was used as the enzyme source. Among the synthesized analogs, 8-thiobutyladenosine 5'-triphosphate (8-BuS-ATP) was found to be the most effective nonhydrolyzable competitive inhibitor, with an estimated K_i of 10 μM . This nonhydrolyzable analog did not exert any P2X-receptor-mediated effect on endothelium-denuded blood vessels, from the guinea pig mesenteric bed. In agreement with this observation, infusion of the analog did not cause any significant blood pressure variations of the precontracted vessel. Because in previous studies on isolated turkey erythrocytes and rat astrocytes 8-BuS-ATP was not able to trigger any P2Y1-receptor-mediated effect, it therefore appears that this **NTPDase** inhibitor does not interfere with purinergic receptors.

IT 284040-53-5 284040-54-6


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)

(synthesis and biochem. and pharmacol. characterizations of novel inhibitors of nucleoside triphosphate diphosphohydrolases)

RN 284040-53-5 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(hexylthio)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 284040-54-6 CAPLUS

CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(butylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 81609-35-0P 284040-51-3P 284040-52-4P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(synthesis and biochem. and pharmacol. characterizations of novel inhibitors of nucleoside triphosphate diphosphohydrolases)


RN 81609-35-0 CAPLUS
CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(ethylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 284040-51-3 CAPLUS
CN Adenosine 5'-(tetrahydrogen triphosphate), 8-(cycloheptylthio)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 284040-52-4 CAPLUS
CN Adenosine 5'-(tetrahydrogen triphosphate), 8-[(2,2-dimethylpropyl)thio]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 61 THERE ARE 61 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT